

NOVEL THERAPEUTIC MOLECULES

FIELD OF THE INVENTION

- 5 The present invention relates generally to novel molecules capable of, *inter alia*, modulating apoptosis in mammalian cells and to genetic sequences encoding same. More particularly, the present invention relates to a novel member of the Bcl-2 family of proteins, referred to herein as "Bim", and to genetic sequences encoding same. The molecules of the present invention are useful, for example, in therapy, diagnosis, antibody
- 10 generation and as a screening tool for therapeutic agents capable of modulating physiological cell death or survival and/or modulating cell cycle entry.

BACKGROUND OF THE INVENTION

- 15 Bibliographic details of the publications referred to by author in this specification are collected at the end of the description. Sequence Identity Numbers (SEQ ID NOs.) for the nucleotide and amino acid sequences referred to in the specification are defined following the bibliography. A summary of the SEQ ID NOs. is provided before the Examples.
- 20 Apoptosis, the physiologic and genetically modulated process of cell death, is of central importance for modelling tissues and maintaining homeostasis in multicellular organisms (Kerr *et al.*, 1972; Jacobson *et al.*, 1997). Great progress is being made towards understanding the biochemistry underlying this intrinsic suicide program. The cellular apoptotic effector molecules include a set of cysteine proteinases, termed caspases, that
- 25 degrade critical cellular substrates (Nicholson and Thornberry, 1997). The regulatory machinery that governs the activation of the caspases is less well understood. However a family of proteins of which Bcl-2 is the prototypic molecule (and is referred to as the Bcl-2 family of proteins) plays a central role (Jacobson, 1997; Reed, 1997; Kroemer, 1997).
- 30 Bcl-2 was the first intracellular regulator of apoptosis to be identified (Vaux *et al.*, 1988) and high levels enhance cell survival under diverse cytotoxic conditions. Other cellular

- 2 -

- homologues, such as Bcl-x_L (Boise *et al.*, 1993) and Bcl-w (Gibson *et al.*, 1996), also enhance cell survival, as do more distantly related viral homologues, such as the adenovirus E1B 19K protein (White *et al.*, 1992) and Epstein-Barr virus BHRF-1 (Henderson *et al.*, 1993). However, the family also includes members such as Bax
- 5 (Oltvai *et al.*, 1993) and Bak (Chittenden *et al.*, 1995; Kiefer *et al.*, 1995; Farrow *et al.*, 1995), which antagonise the activity of the pro-survival proteins and provoke apoptosis when expressed at high concentrations. The relative concentrations of the opposing sub-family members may determine whether the cell lives or dies (Oltvai *et al.*, 1993).
- 10 The homology between members of the Bcl-2 family is greatest within four small regions, designated Bcl-2 Homology (BH) regions (Yin *et al.*, 1994; Borner *et al.*, 1994; Chittenden *et al.*, 1995; Gibson *et al.*, 1996; Zha *et al.*, 1996). The N-terminal BH4 domain is restricted to some antagonists of apoptosis, while BH1, BH2 and BH3 can be found in both sub-families (reviewed by Kroemer, 1997). In the tertiary structure
- 15 determined for Bcl-x_L, the BH1, BH2 and BH3 domains form an elongated hydrophobic cleft on the surface of the molecule, stabilised by the BH4 amphipathic helix (Muchmore *et al.*, 1996; Sattler *et al.*, 1997). Most members of the Bcl-2 family contain a C-terminal hydrophobic region, which appears to be important for their localisation to intracytoplasmic membranes (reviewed by Kroemer, 1997).
- 20 Protein interactions are an important feature of the Bcl-2 family. Interaction between the pro-survival and pro-apoptotic proteins, such as Bcl-2 with Bax or Bak, requires the BH1 and BH2 domains of the former (Yin *et al.*, 1994; Sedlak *et al.*, 1995; Hanada *et al.*, 1995) and the BH3 domain of the latter (Chittenden *et al.*, 1995; Zha *et al.*, 1996). BH3
- 25 peptides bind to the hydrophobic cleft of Bcl-x_L formed by BH1, BH2 and BH3 (Sattler *et al.*, 1997). Although mutagenesis of Bcl-2 and Bcl-x_L initially suggested that their ability to inhibit cell death required binding to a pro-apoptotic family member (Yin *et al.*, 1994), Bcl-x_L mutants have been identified that do not bind Bax or Bak but still block apoptosis (Cheng *et al.*, 1996).

- 3 -

An additional group of pro-apoptotic proteins has recently been described. Bik/Nbk (Boyd *et al.*, 1995; Zha *et al.*, 1996), Bid (Wang *et al.*, 1996) and Hrk (Inohara *et al.*, 1997). The only feature they share in common with each other, or the Bcl-2 family, is the small (9 amino acid) BH3 domain. This region is essential for the ability of these proteins to promote cell death.

In work leading up to the present invention, the inventors have identified a novel member of the Bcl-2 family, designated herein "Bim". In accordance with the present invention, Bim induces cell death and acts as a "death-ligand" for certain members of the pro-survival Bcl-2 family. The identification of this new gene permits the identification and rational design of a range of products for use in therapy, diagnosis, antibody generation and involving modulation of physiological cell death. These therapeutic molecules may act as either antagonists or agonists of Bim's function and will be useful in cancer autoimmune or degenerative disease therapy.

15

SUMMARY OF THE INVENTION

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

Specific mutations in an amino acid sequence are represented herein as "X₁nX₂" where X₁ is the original amino acid residue before mutation, n is the residue number and X₂ is the mutant amino acid. Reference to X_n is a reference to a particular amino acid in an amino acid sequence where X is the amino acid and n is the residue number. The abbreviation X may be to the three letter or single letter amino acid code.

One aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide having one or more of the identifying characteristics of Bim or a derivative or homologue thereof.

- 4 -

Another aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence encoding or complementary to a sequence encoding an amino acid sequence substantially as set forth in one of SEQ ID NO: 2, 4, or 6 or a derivative or homologue thereof or having at least about 45% or greater similarity to one or more of
5 SEQ ID NO: 2, 4, or 6, or a derivative or homologue thereof.

Another aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence encoding or complementary to a sequence encoding an amino acid sequence substantially as set forth in one of SEQ ID NO: 8 or 10 or a derivative or
10 homologue thereof or having at least about 45% or greater similarity to one or more of SEQ ID NO: 8 or 10 or a derivative or homologue thereof.

Yet another aspect of the present invention contemplates a nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or
15 5 or a derivative or homologue thereof capable of hybridising to one of SEQ ID NO: 1, 3, or 5 under low stringency conditions at 42°C and which encodes an amino acid sequence corresponding to an amino acid sequence set forth in one of SEQ ID NO: 2, 4 or 6 or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

20 Still yet another aspect of the present invention contemplates a nucleic acid molecule comprising a sequence of nucleotides substantially as set forth in SEQ ID NO: 1, 3, or 5.

Still another aspect of the present invention contemplates a nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9
25 or a derivative of homologue thereof capable of hybridising to one of SEQ ID NO: 7 or 9 under low stringency conditions at 42°C and which encodes an amino acid sequence corresponding to an amino acid sequence set forth in one of SEQ ID NO: 8 or 10 or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

30 A further aspect of the present invention contemplates a nucleic acid molecule comprising

- 5 -

a sequence of nucleotides substantially as set forth in SEQ ID NO: 7 or 9.

Another further aspect of the present invention is directed to an isolated nucleic acid molecule encoding *Bim* or a derivative thereof, said nucleic acid molecule selected from the list consisting of:

- (i) A nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence set forth in one of SEQ ID NO: 2, 4, or 6 or a derivative or homologue thereof or having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- (ii) A nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence set forth in one of SEQ ID NO: 8 or 10 or a derivative or homologue or having at least about 45% similarity to one of SEQ ID NO: 8 or 10.
- (iii) A nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or 5 or a derivative or homologue thereof.
- (iv) A nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9 or a derivative or homologue thereof.
- (v) A nucleic acid molecule capable of hybridising under low stringency conditions at 42°C to the nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or 5 a derivative or homologue and encoding an amino acid sequence corresponding to an amino acid sequence as set forth in one of SEQ ID NO: 2, 4 or 6 a derivative or homologue or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- (vi) A nucleic acid molecule capable of hybridising under low stringency conditions at 42°C to the nucleotide sequence substantially as set forth in one of SEQ ID NO: 7

- 6 -

or 9 a derivative or homologue and encoding an amino acid sequence corresponding to an amino acid sequence as set forth in one of SEQ ID NO: 8 or 10 a derivative or homologue or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

5

- (vii) A nucleic acid molecule capable of hybridising to the nucleic acid molecule of paragraphs (i) or (iii) or (v) under low stringency conditions at 42°C and encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

10

- (viii) A nucleic acid molecule capable of hybridising to the nucleic acid molecule of paragraphs (ii) or (iv) or (vi) under low stringency conditions at 42°C and encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

15

- (ix) A derivative or mammalian homologue of the nucleic acid molecule of paragraphs (i) or (ii) or (iii) or (iv) or (v) or (vi) or (vii) or (viii).

Yet another further aspect of the present invention is directed to an isolated polypeptide
20 selected from the list consisting of:

- (i) A polypeptide having an amino acid sequence substantially as set forth in one of SEQ ID NO: 2, 4, or 6 or derivative or homologue thereof or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

25

- (ii) A polypeptide having an amino acid sequence substantially as set forth in one of SEQ ID NO: 8 or 10 a derivative or homologue or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

- 30 (iii) A polypeptide encoded by a nucleotide sequence substantially as set forth in one of

- 7 -

SEQ ID NO: 1, 3, or 5 or derivative or homologue thereof or a sequence encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

- 5 (iv) A polypeptide encoded by a nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9 or derivative or homologue thereof or a sequence encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.
- 10 (v) A polypeptide encoded by a nucleic acid molecule capable of hybridising to the nucleotide sequence as set forth in one of SEQ ID NO: 1, 3, or 5 or derivative or homologue thereof under low stringency conditions at 42°C and which encodes an amino acid sequence substantially as set forth in SEQ ID NO: 2, 4, or 6 or
- 15 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- (vi) A polypeptide encoded by a nucleic acid molecule capable of hybridising to the nucleotide sequence as set forth in one of SEQ ID NO: 7 or 9 or derivative or homologue thereof under low stringency conditions at 42°C and which encodes an
- 20 amino acid sequence substantially as set forth in SEQ ID NO: 8 or 10 or derivative or homologue thereof or an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.
- (vii) A polypeptide as defined in paragraphs (i) or (ii) or (iii) or (iv) or (v) or (vi) in
- 25 homodimeric form.
- (viii) A polypeptide as defined in paragraphs (i) or (ii) or (iii) or (iv) or (v) or (vi) in heterodimeric form.

30 Accordingly, a related aspect of the present invention is directed to a variant of an isolated

- 8 -

Bim nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion to the polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

5

Accordingly, the present invention is more particularly directed to a variant of an isolated *Bim* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion in the region of the polypeptide encoded by said variant which binds the dynein light chain
10 wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

Even more preferably, the present invention is directed to a variant of an isolated murine or human *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations in said
15 nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion in the region defined by amino acid residue numbers 42 to 71 of the polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

20 In another preferred embodiment the present invention is directed to a variant of an isolated murine *Bim_{EL}* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion in the region defined by amino acid residue numbers 42 to 127 of the polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or
25 otherwise associate with a dynein light chain.

In yet another preferred embodiment the present invention is directed to a variant of an isolated human *Bim_{EL}* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution
30 and/or deletion in the region defined by amino acid residue numbers 42 to 131 of the

- 9 -

polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

Accordingly, a preferred embodiment of the present invention is directed to a variant of an
5 isolated human or murine *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitution D51G of the polypeptide encoded by said mutated nucleic acid molecule.

Another preferred embodiment of the present invention is directed to a variant of an
10 isolated human or murine *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitution S53P of the polypeptide encoded by said mutated nucleic acid molecule.

In another preferred embodiment the present invention provides a variant of an isolated
15 human or murine *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitution T54A of the polypeptide encoded by said mutated nucleic acid molecule.

In yet another preferred embodiment the present invention provides a variant of an isolated
20 *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitutions T54I and N65S of the polypeptide encoded by said mutated nucleic acid molecule.

Accordingly, the present invention is directed to a variant of an isolated *Bim* polypeptide
25 comprising at least one amino acid addition, substitution and/or deletion wherein said polypeptide cannot bind, couple or otherwise associate with the dynein light chain.

Another aspect of the present invention contemplates a method for modulating expression of *Bim* in a mammal, said method comprising administering to said mammal a modulating
30 effective amount of an agent for a time and under conditions sufficient to up-regulate or

- 10 -

down-regulate or otherwise modulate expression of *Bim*.

Yet another aspect of the present invention contemplates a method of modulating activity of *Bim* in a mammal, said method comprising administering to said mammal a modulating
5 effective amount of an agent for a time and under conditions sufficient to increase or decrease *Bim* activity.

Still another aspect of the present invention contemplates a method of modulating apoptosis in a mammal said method comprising administering to said mammal an effective
10 amount of an agent for a time and under conditions sufficient to modulate the expression of a nucleotide sequence encoding *Bim*.

Yet another aspect of the present invention contemplates a method of modulating apoptosis in a mammal said method comprising administering to said mammal an effective amount
15 of an agent for a time and under conditions sufficient to modulate the activity of *Bim*.

Still another aspect of the present invention contemplates a method of modulating apoptosis in a mammal said method comprising administering to said mammal an effective amount of *Bim* or *Bim* or derivative thereof.

20

Accordingly, another aspect of the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate the expression of *Bim* wherein said modulation results in modulation of apoptosis.

25

In another aspect the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate the activity of *Bim* wherein said modulation results in modulation of apoptosis.

30

- 11 -

In another aspect the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of Bim or derivative thereof for a time and under conditions sufficient to modulate apoptosis.

- 5 Yet another aspect the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of *Bim* or derivative thereof for a time and under conditions sufficient to modulate apoptosis.

- 10 In yet another aspect the present invention relates to the use of an agent capable of modulating the expression of *Bim* in the manufacture of a medicament for the modulation of apoptosis.

- 15 Still yet another aspect of the present invention relates to the use of an agent capable of modulating the expression of Bim in the manufacture of a medicament for the modulation of apoptosis.

A further aspect of the present invention relates to the use of Bim or *Bim* or derivative thereof in the manufacture of a medicament for the modulation of apoptosis.

- 20 Another further aspect of the present invention relates to agents for use in modulating *Bim* expression wherein modulating expression of said *Bim* modulates apoptosis.

- 25 Yet another further aspect of the present invention relates to agents for use in modulating Bim expression wherein modulating expression of said Bim modulates apoptosis.

Still yet another further aspect of the present invention relates to Bim or *Bim* or derivative thereof for use in modulating apoptosis.

- 30 Another aspect of the present invention contemplates a pharmaceutical composition comprising *Bim*, Bim or derivative thereof or an agent capable of modulating *Bim*

- 12 -

expression or Bim activity together with one or more pharmaceutically acceptable carriers and/or diluents.

Yet another aspect of the present invention is directed to an immunointeractive molecule
5 comprising an antigen binding portion having specificity for Bim or *Bim* or derivative thereof.

Still another aspect of the present invention contemplates a monoclonal antibody having specificity for Bim or *Bim* or derivative thereof.

10

Still yet another aspect of the present invention provides a monoclonal antibody having specificity for Bim_L.

A further aspect of the present invention provides a method of detecting an
15 immunointeractive molecule, in a sample, specific for a protein of interest produced by a cell said method comprising contacting the sample to be tested with a population of cells comprising a defined ratio of cells producing the protein of interest and cells not producing the protein of interest for a time and under conditions sufficient for immunointeractive molecules, if present in said sample, to interact with said protein of
20 interest and the subjecting said immunointeractive molecule-protein complex to detecting means.

Another further aspect of the present invention contemplates a method for detecting Bim in a biological sample from a subject said method comprising contacting said biological
25 sample with an immunointeractive molecule as hereinbefore defined specific for Bim or its derivatives thereof for a time and under conditions sufficient for an immunointeractive molecule-Bim complex to form, and then detecting said complex.

Yet another further aspect of the present invention contemplates a method for detecting
30 *Bim* in a biological sample from a subject said method comprising contacting said

- 13 -

biological sample with an immunointeractive molecule as hereinbefore defined specific for *Bim* or its derivatives thereof for a time and under conditions sufficient for an immunointeractive molecule-*Bim* complex to form, and then detecting said complex.

- 5 Single and three letter abbreviations used throughout the specification are defined in Table 1.

TABLE 1
Single and three letter amino acid abbreviations

	Amino Acid	Three-letter Abbreviation	One-letter Symbol
5			
	Alanine	Ala	A
	Arginine	Arg	R
	Asparagine	Asn	N
10	Aspartic acid	Asp	D
	Cysteine	Cys	C
	Glutamine	Gln	Q
	Glutamic acid	Glu	E
	Glycine	Gly	G
15	Histidine	His	H
	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K
	Methionine	Met	M
20	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
	Threonine	Thr	T
	Tryptophan	Trp	W
25	Tyrosine	Tyr	Y
	Valine	Val	V
	Any residue	Xaa	X

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of the isolation of cDNAs encoding three isoforms of Bim. (A) Open reading frames of five independent clones isolated by screening a cDNA expression library with recombinant Bcl-2 protein. Dotted lines indicate putative splices and arrows indicate PCR primers spanning the splice sites. (B) Relationship of the three Bim isoforms. The black box denotes the BH3 homology region and the hatched box the predicted hydrophobic region. Regions specific to the larger splice variants are shaded. (C) Sequence alignment of the mouse and human Bim_{EL} polypeptide sequences using the GCG "BESTFIT" program; identical residues appear on a dark background. The BH3 homology region and the C-terminal hydrophobic region predicted by the Kyte-Doolittle algorithm are boxed. Arrows indicate residues present only in the longer isoforms. Since the nucleotide sequences of the mouse and human cDNAs diverged 5' of the predicted initiating ATG and there are stop codons in all three reading frames upstream of the human open reading frame, that start codon is likely to be correct.

Figure 2 is a photographic representation of the expression of *bim* RNA in haematopoietic cell lines. Northern blot analysis of polyA⁺ RNA, using a mouse *bim* cDNA probe. The RNAs were derived from the following mouse lines: T lymphomas KO52DA20 (lanes 1 to 5), WEHI 703 (lane 6), WEHI 707 (lane 7) and WEHI 7.1 (lane 8); B lymphomas CH1 (lanes 9, 10) and WEHI 231 (lanes 11, 12); pre-B lymphoma WEHI 415 (lane 13); T hybridoma B6.2.16 BW2 (lanes 14, 15); myeloid progenitor FDC-P1 (lane 16). Those lines that harbour a *bcl-2* expression vector or transgene are indicated. Certain RNAs were isolated from cells exposed to cytotoxic conditions: 1 μ M dexamethasone (14 hr, lanes 2 and 4; 24 hr, lane 5); γ -irradiation (10 Gy) (lane 5). Samples from a single autoradiograph have been rearranged in order electronically.

Figure 3 is a photographic representation of the localisation of Bim protein to intracellular membranes. (A) L929 fibroblasts transiently transfected with EE-tagged Bim_L were fixed, permeabilised and stained with the anti-EE antibody; fluorescence was visualised by confocal

microscopy. (B) and (C) L929 cells stably co-expressing human Bcl-2 and EE-tagged Bim_L were stained with anti-human Bcl-2 antibody (B) or anti-EE antibody (C). (D) Images from the staining with anti-EE (B) and anti-Bcl-2 (C) were superimposed; co-localisation is indicated by (*colour) staining.

5

Figure 4 is a graphical representation demonstrating that Bim induces apoptosis and can be inhibited by p35 and Bcl-2 but not CrmA. (A) Flow cytometric DNA analysis (see Materials and Methods) of 293T cells transfected 24 h previously with *EE-bim_L* plasmid (0.5 µg). (B) Kinetics of apoptosis elicited by *EF-bim_L* plasmid (0.5 µg), assessed as in 10 A. (C) Cell viability 48 h after transfection with 0.1, 0.2 or 0.5 µg of *EE-bim_L* plasmid alone (black bars) or together with 0.5 µg of wild-type or mutant *p35* or *crmA* plasmid (grey bars). (D) Cell viability 48 h after transfection with 0.1, 0.2 or 0.5 µg of *EE-bim_L* plasmid together with 0.5 µg of the indicated wt or mutant *bcl-2* plasmids. C and D show the percentage of viable Bim-expressing cells, determined by DNA FACS analysis, 15 as in A, and are the mean \pm SD of 3 or more independent experiments.

Figure 5 is a graphical representation demonstrating that Bim antagonises the anti-apoptotic activity of Bcl-2 in a dose-dependent fashion. (A) Immunofluorescence staining of cloned FDC-P1 cell lines stably expressing Bcl-2 alone (dashed line) or co-expressing 20 Bcl-2 and varying levels of EE-Bim_L (solid lines). (B) Viability of these clones when cultured in the absence of IL-3 or after exposure to γ -irradiation (10 Gy). Cell viability was assessed by vital dye exclusion; data shown are means \pm SD of at least 3 experiments and are representative of results obtained with at least 3 independent lines of each genotype.

25

Figure 6 is a graphical representation of a comparison of the activity of the three Bim isoforms. (A) Immunofluorescence staining of cloned FDC-P1 lines expressing Bcl-2 alone (dotted) or Bcl-2 plus EE-tagged Bim_L, Bim_{EL} or Bim_S (solid lines). (B) Association of EE-tagged Bim_S, Bim_L and Bim_{EL} with Bcl-2 demonstrated by anti-EE 30 immunoblots of immunoprecipitates prepared with anti-human Bcl-2 monoclonal antibody

from FDC-P1 cells expressing the indicated proteins. The 25 kD protein is non-specific.

(C) Effect of Bim isoforms on viability of FDC-P1 cells expressing Bcl-2, after removal of growth factor or exposure to irradiation.

- 5 **Figure 7** is a graphical representation demonstrating that Bim binds to and antagonises Bcl-x_L or Bcl-w but not E1B19K. (A) Lysates of ³⁵S-labelled 293T cells transiently co-transfected with the plasmids encoding the indicated proteins were immunoprecipitated with anti-EE antibody, and the EE-Bim_L-containing complexes were fractionated by SDS-PAGE. (B) Lysates from parental 293T cells or 293T cells co-expressing EE-tagged
- 10 Bim_L and FLAG-tagged Bcl-x_L, Bcl-w or E1B19K were immunoblotted directly or after immunoprecipitation, as indicated. (C, D) 293T cells were transiently transfected with a vector control (unfilled bar) or with 0.1, 0.2 or 0.5 µg of EE-Bim_L plasmid, either alone (black bars) or together with 0.5 µg of plasmids encoding wt or mutant Bcl-x_L (C); Bcl-w or E1B19K protein (D) (grey bars). The flow cytometric analysis was as described in the
- 15 legend to Figure 4. Data shown are means ± SD of 3 or more independent experiments.

Figure 8 is a graphical representation demonstrating that the BH3 homology region of Bim is required for binding to and inhibiting Bcl-2. (A) Immunofluorescence staining of cloned FDC-P1 lines expressing Bcl-2 alone (dotted) or with EE-Bim_L or EE-Bim ΔBH3

20 (solid line), and of EE-Bim ΔBH3 in the parental FDC-P1 cells (broken line). (B) Immunoblot showing that Bcl-2 associates with wild-type Bim_L but not the ΔBH3 mutant. (C) Viability of FDC-P1 clones expressing the indicated proteins (see A) was assessed by vital dye exclusion. Data shown are means ± SD of at least 3 experiments and are representative of results obtained with at least 3 independent lines of each genotype.


25

Figure 9 is a diagrammatic representation of the BH3 homology regions in the Bcl-2 family. (A) Amino acid sequences of the human proteins were aligned with the modified method of Feng and Doolittle used by the GCG "PILEUP" program (Feng and Doolittle, 1987). Residues that are identical or very similar (K & R; D & E; V & I; M & L) in > 8

30 of the 11 proteins are shaded in dark grey, while less conserved residues (present in

Ant
B1

- 18 -


 > 5/11 proteins) are shaded in light grey. (B) Short stretch of amino acid homology between Bim and *C. elegans* Ced-4; this region overlaps with the BH3 region of Bim, indicated by the box.

5 **Figure 10** is a graphical representation of the detection of Bim Specific Antibodies.

Stably transfected FDC-P1/Bcl-2 and FDC-P1/Bcl-2/EE-Bim cells were mixed 1:1, fixed, permeabilized and stained with anti-EE antibodies (A, positive control) or with hybridoma supernatants from the fusion (B-F). (B) shows a typical negative clone, (C) an antibody that is not specific to Bim but recognizes an epitope that is present in both FDC-P1/Bcl-2
 10 and FDC-P1/Bcl-2/EE-Bim cells, and D-F show staining by the anti-Bim antibodies 4E4 (D), 5E5 (E) and 9F5 (F). Staining was visualized by either FITC-conjugated goat anti-mouse IgG antibodies (A) or FITC-conjugated goat anti-rat IgG antibodies (B-F), and analyzed by flow cytometry. Supernatants with anti-Bim reactivity (D-F) produced a double peak: background staining of the FDC-P1/Bcl-2 cells (lower intensity peak) and
 15 specific Bim staining of the FDC-P1/Bcl-2/EE-Bim cells (higher intensity peak).

Figure 11 is a graphical representation of the determination of the Ig isotypes of the Bim-reactive monoclonal antibodies. A 1:1 mixture of stably transfected FDC-P1/Bcl-2 and FDC-P1/Bcl-2/EE-Bim cells was fixed, permeabilized and stained with anti-Bim antibody
 20 (clone 9F5), followed by either of the biotinylated mouse anti-rat Ig isotype specific monoclonal antibodies: anti-rat IgG1 (A), anti-rat IgG2a (B), anti-rat IgG2b (C), a control antibody (D) or directly with anti-rat Ig κ light chain conjugated to FITC (E and F [negative control]). In the case of staining with biotinylated antibodies, FITC-coupled streptavidin was used as the tertiary reagent. The double peaks (B and E) indicate that
 25 9F5 is an antibody of the IgG2a/ κ isotype.

Figure 12 is a photographic representation of anti-Bim monoclonal antibody detecting Bim by Western blotting. Expression of EE-Bim_L in FDC-P1/Bcl-2/EE-Bim_L cell lysates (10³-10⁶ cells) was analyzed by Western blotting using the anti-Bim antibody (9F5) and
 30 goat anti-rat Ig conjugated to HRP as a secondary reagent and detection by enhanced

- 19 -

chemiluminescence. Lysates from FDC-P1/Bcl-2 cells were used as negative controls. A specific band of ~23kD, corresponding to EE Bim_L, was only detected in FDC-P1/Bcl-2/EE-Bim_L lysates and could not be detected in lysates from fewer than 5 x 10⁴ cells. The upper band in the lane which contains lysate from 10⁶

5 FDC-P1/Bcl-2/EE-Bim cells is an artefact of overloading which led to retention of some Bim protein during electrophoresis.

Figure 13 is a photographic representation of anti-Bim monoclonal antibody detecting Bim by immunoprecipitation. Equivalent lysates from metabolically labelled 293T cells (lanes 2, 4 and 6) or 293T cells transiently transfected with a FLAG-Bim_L expression construct (lanes 1, 3 and 5) were immunoprecipitated using the anti-FLAG antibody M2 (lanes 1 and 2), the anti-Bim antibody 5E5 (lanes 3 and 4) or an isotype-matched control antibody to mouse CD4 (lanes 5 and 6). Both the anti-FLAG antibody and the anti-Bim 5E5 antibody immunoprecipitated a 23 kD protein that corresponded with the expected mobility of FLAG-Bim_L.

Figure 14 is a schematic representation of the gene targeting vector utilised in making Bim knockout mice.

- 20 -

DETAILED DESCRIPTION OF THE INVENTION

Accordingly, one aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence encoding or complementary to a sequence encoding an amino acid sequence substantially as set forth in one of SEQ ID NO: 2, 4, or 6 or a derivative or homologue thereof or having at least about 45% or greater similarity to one or more of SEQ ID NO: 2, 4, or 6, or a derivative or homologue thereof.

Another aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence encoding or complementary to a sequence encoding an amino acid sequence substantially as set forth in one of SEQ ID NO: 8 or 10 or a derivative or homologue thereof or having at least about 45% or greater similarity to one or more of SEQ ID NO: 8 or 10 or a derivative or homologue thereof.

15 The term "similarity" as used herein includes exact identity between compared sequences at the nucleotide or amino acid level. Where there is non-identity at the nucleotide level, "similarity" includes differences between sequences which result in different amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. Where there is non-identity at the amino acid level, "similarity" includes amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. In a particular preferred embodiment, nucleotide and sequence comparisons are made at the level of identity rather than similarity. Any number of programs are available to compare nucleotide and amino acid sequences. Preferred programs have regard to an appropriate alignment. One such program is Gap which considers all possible alignment and gap positions and creates an alignment with the largest number of matched bases and the fewest gaps. Gap uses the alignment method of Needleman and Wunsch. Gap reads a scoring matrix that contains values for every possible GCG symbol match. GAP is available on ANGIS (Australian National Genomic Information Service) at website <http://mell1.angis.org.au..>

- 21 -

Another aspect of the present invention contemplates a nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or 5 or a derivative or homologue thereof capable of hybridising to one of SEQ ID NO: 1, 3, or 5 under low stringency conditions at 42°C and which encodes an amino acid sequence
5 corresponding to an amino acid sequence set forth in one of SEQ ID NO: 2, 4 or 6 or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

More particularly the present invention contemplates a nucleic acid molecule comprising a sequence of nucleotides substantially as set forth in SEQ ID NO: 1, 3, or 5.

10

Another aspect of the present invention contemplates a nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9 or a derivative of homologue thereof capable of hybridising to one of SEQ ID NO: 7 or 9 under low stringency conditions at 42°C and which encodes an amino acid sequence corresponding to
15 an amino acid sequence set forth in one of SEQ ID NO: 8 or 10 or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

More particularly the present invention contemplates a nucleic acid molecule comprising a sequence of nucleotides substantially as set forth in SEQ ID NO: 7 or 9.

20

Reference herein to a low stringency at 42°C includes and encompasses from at least about 1% v/v to at least about 15% v/v formamide and from at least about 1M to at least about 2M salt for hybridisation, and at least about 1M to at least about 2M salt for washing conditions. Alternative stringency conditions may be applied where necessary,
25 such as medium stringency, which includes and encompasses from at least about 16% v/v to at least about 30% v/v formamide and from at least about 0.5M to at least about 0.9M salt for hybridisation, and at least about 0.5M to at least about 0.9M salt for washing conditions, or high stringency, which includes and encompasses from at least about 31% v/v to at least about 50% v/v formamide and from at least about 0.01M to at least about
30 0.15M salt for hybridisation, and at least about 0.01M to at least about 0.15M salt for

- 22 -

washing conditions. In general, washing is carried out at $T_m = 69.3 + 0.41 (G + C)\%[19] = -12^\circ\text{C}$. However, the T_m of a duplex DNA decreases by 1°C with every increase of 1% in the number of mismatched based pairs (20).

5 The nucleic acid molecule according to this aspect of the present invention corresponds herein to "*Bim*". This gene has been determined in accordance with the present invention to induce apoptosis. The product of the *Bim* gene is referred to herein as "Bim" without limiting this invention in any way, murine *Bim* has been mapped to murine chromosome 2 at bands F3-G and human *Bim* has been mapped to the syntenic region on chromosome 2
10 at bands 2q12-2q13. Bim is known as a "BH3-only" protein since the only Bcl-2 homology region which it encompasses is BH3. It thereby forms a novel member of a Bcl-2 related BH3-only pro-apoptotic group which also comprises, for example, Bik/Nbk, Bid and Hrk. However, Bim is the only BH3-only protein for which splice variants exist, thereby resulting in the expression of a variety of isoforms. Bim_S, Bim_L and Bim_{EL} are
15 examples of three said isoforms which differ in both size and potency of functional activity. Murine Bim_S, Bim_L and Bim_{EL} are defined by the amino acid sequences set forth in SEQ ID NO: 2, 4 and 6, respectively and human Bim_L and Bim_{EL} are defined by the amino acid sequences set forth in SEQ ID NO: 8 and 10, respectively. The cDNA
20 nucleotide sequences for murine Bim_S, Bim_L and Bim_{EL} are defined by the nucleotide sequences set forth in SEQ ID NO: 1, 3 and 5, respectively and human Bim_L and Bim_{EL} are defined by the nucleotide sequences set forth in SEQ ID NO: 7 and 9, respectively.

The nucleic acid molecule encoding *Bim* is preferably a sequence of deoxyribonucleic acids such as cDNA sequence, an mRNA sequence or a genomic sequence. A genomic
25 sequence may also comprise exons and introns. A genomic sequence may also include a promoter region or other regulatory region.

Reference hereinafter to "Bim" and "*Bim*" should be understood as a reference to all forms of Bim and *Bim*, respectively, including, by way of example, the three peptide and cDNA
30 isoforms of Bim_S, Bim_L and Bim_{EL} which have been identified as arising from alternative

- 23 -

splicing of mRNA and the *Bim* gene. Reference hereinafter to *Bim* and *Bim* in the absence of a reference to its derivatives should be understood to include reference to its derivatives thereof.

- 5 The protein and/or gene is preferably from a human, primate, livestock animal (eg. sheep, pig, cow, horse, donkey) laboratory test animal (eg. mouse, rat, rabbit, guinea pig) companion animal (eg. dog, cat), captive wild animal (eg. fox, kangaroo, deer), aves (eg. chicken, geese, duck, emu, ostrich), reptile or fish.
- 10 Derivatives include fragments (such as peptides), parts, portions, chemical equivalents, mutants, homologues or mimetics from natural, synthetic or recombinant sources including fusion proteins. Derivatives may be derived from insertion, deletion or substitution of amino acids. Amino acid insertional derivatives include amino and/or carboxylic terminal fusions as well as intrasequence insertions of single or multiple amino
15 acids. Insertional amino acid sequence variants are those in which one or more amino acid residues are introduced into a predetermined site in the protein although random insertion is also possible with suitable screening of the resulting product. Deletional variants are characterized by the removal of one or more amino acids from the sequence. Substitutional amino acid variants are those in which at least one residue in the sequence
20 has been removed and a different residue inserted in its place. Additions to amino acid sequences including fusions with other peptides, polypeptides or proteins. Mutants should be understood to include, but is not limited to, the specific *Bim* or *Bim* mutant molecules described herein. Derivatives include, for example, peptides derived from the BH3 region, from the dynein binding region or from a site of phosphorylation. Peptides
25 include, for example, molecules comprising at least 4 contiguous amino acids corresponding to at least 4 contiguous amino acids of *Bim* as herein defined. Use of the term "polypeptides" herein should be understood to encompass peptides, polypeptides and proteins.
- 30 The derivatives of *Bim* include fragments having particular epitopes or parts of the entire *Bim* protein fused to peptides, polypeptides or other proteinaceous or non-proteinaceous molecules. For example, *Bim* or derivative thereof may be fused to a molecule to

- 24 -

facilitate its entry into a cell. Analogues of Bim contemplated herein include, but are not limited to, modification to side chains, incorporating of unnatural amino acids and/or their derivatives during peptide, polypeptide or protein synthesis and the use of crosslinkers and other methods which impose conformational constraints on the proteinaceous molecules or
5 their analogues. Derivatives of nucleic acid sequences may similarly be derived from single or multiple nucleotide substitutions, deletions and/or additions including fusion with other nucleic acid molecules. The derivatives of the nucleic acid molecules of the present invention include oligonucleotides, PCR primers, antisense molecules, molecules suitable for use in cosuppression and fusion of nucleic acid molecules.

10

Examples of side chain modifications contemplated by the present invention include modifications of amino groups such as by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH_4 ; amidination with methylacetimidate; acylation with acetic anhydride; carbamoylation of amino groups with cyanate;
15 trinitrobenzylation of amino groups with 2, 4, 6-trinitrobenzene sulphonic acid (TNBS); acylation of amino groups with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation of lysine with pyridoxal-5-phosphate followed by reduction with NaBH_4 .

The guanidine group of arginine residues may be modified by the formation of
20 heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

The carboxyl group may be modified by carbodiimide activation *via* O-acylisourea formation followed by subsequent derivitisation, for example, to a corresponding amide.

25

Sulphydryl groups may be modified by methods such as carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulphides with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted maleimide; formation of mercurial derivatives using 4-
30 chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-

- 25 -

chloromercuri-4-nitrophenol and other mercurials; carbamoylation with cyanate at alkaline pH.

Tryptophan residues may be modified by, for example, oxidation with N-

- 5 bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues on the other hand, may be altered by nitration with tetranitromethane to form a 3-nitrotyrosine derivative.

Modification of the imidazole ring of a histidine residue may be accomplished by

- 10 alkylation with iodoacetic acid derivatives or N-carboethoxylation with diethylpyrocarbonate.

Examples of incorporating unnatural amino acids and derivatives during protein synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-

- 15 hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl alanine and/or D-isomers of amino acids. A list of unnatural amino acid contemplated herein is shown in Table 2.

TABLE 2

	Non-conventional amino acid	Code	Non-conventional amino acid	Code
5				
	α -aminobutyric acid	Abu	L-N-methylalanine	Nmala
	α -amino- α -methylbutyrate	Mgab	L-N-methylarginine	Nmarg
	aminocyclopropane- carboxylate	Cpro	L-N-methylasparagine	Nmasn
			L-N-methylaspartic acid	Nmasp
10	aminoisobutyric acid	Aib	L-N-methylcysteine	Nmcys
	aminonorbornyl- carboxylate	Norb	L-N-methylglutamine	Nmgln
			L-N-methylglutamic acid	Nmglu
	cyclohexylalanine		Chexa L-N-methylhistidine	Nmhis
	cyclopentylalanine	Cpen	L-N-methylisoleucine	Nmile
15	D-alanine	Dal	L-N-methylleucine	Nmleu
	D-arginine	Darg	L-N-methyllysine	Nmlys
	D-aspartic acid	Das	L-N-methylmethionine	Nmmet
	D-cysteine	Dcys	L-N-methylnorleucine	Nmnle
	D-glutamine	Dgln	L-N-methylnorvaline	Nmnva
20	D-glutamic acid	Dglu	L-N-methylornithine	Nmorn
	D-histidine	Dhis	L-N-methylphenylalanine	Nmphe
	D-isoleucine	Dile	L-N-methylproline	Nmpro
	D-leucine	Dleu	L-N-methylserine	Nmser
	D-lysine	Dlys	L-N-methylthreonine	Nmthr
25	D-methionine	Dmet	L-N-methyltryptophan	Nmtrp
	D-ornithine	Dorn	L-N-methyltyrosine	Nmtyr
	D-phenylalanine	Dphe	L-N-methylvaline	Nmval
	D-proline	Dpro	L-N-methylethylglycine	Nmetg
	D-serine	Dser	L-N-methyl-t-butylglycine	Nmtbug
30	D-threonine	Dthr	L-norleucine	Nle

	D-tryptophan	Dtrp	L-norvaline	Nva
	D-tyrosine	Dtyr	α -methyl-aminoisobutyrate	Maib
	D-valine	Dval	α -methyl- γ -aminobutyrate	Mgab
	D- α -methylalanine	Dmala	α -methylcyclohexylalanine	Mchexa
5	D- α -methylarginine	Dmarg	α -methylcyclopentylalanine	Mcpen
	D- α -methylassparagine	Dmasn	α -methyl- α -naphthylalanine	Manap
	D- α -methylasspartate	Dmasp	α -methylpenicillamine	Mpen
	D- α -methylcysteine	Dmcys	N-(4-aminobutyl)glycine	Nglu
	D- α -methylglutamine	Dmgln	N-(2-aminoethyl)glycine	Naeg
10	D- α -methylhistidine	Dmhis	N-(3-aminopropyl)glycine	Norn
	D- α -methylisoleucine	Dmile	N-amino- α -methylbutyrate	Nmaabu
	D- α -methyllleucine	Dmleu	α -naphthylalanine	Anap
	D- α -methyllysine	Dmlys	N-benzylglycine	Nphe
	D- α -methylmethionine	Dmmet	N-(2-carbamylethyl)glycine	Ngln
15	D- α -methylornithine	Dmorn	N-(carbamylmethyl)glycine	Nasn
	D- α -methylphenylalanine	Dmphe	N-(2-carboxyethyl)glycine	Nglu
	D- α -methylproline	Dmpro	N-(carboxymethyl)glycine	Nasp
	D- α -methylserine	Dmser	N-cyclobutylglycine	Ncbut
	D- α -methylthreonine	Dmthr	N-cycloheptylglycine	Nchep
20	D- α -methyltryptophan	Dmtrp	N-cyclohexylglycine	Nchex
	D- α -methyltyrosine	Dmtty	N-cyclodecylglycine	Ncdec
	D- α -methylvaline	Dmval	N-cylcododecylglycine	Ncdod
	D-N-methylalanine	Dnmala	N-cyclooctylglycine	Ncoct
	D-N-methylarginine	Dnmarg	N-cyclopropylglycine	Ncpro
25	D-N-methylassparagine	Dnmasn	N-cycloundecylglycine	Ncund
	D-N-methylasspartate	Dnmasp	N-(2,2-diphenylethyl)glycine	Nbhm
	D-N-methylcysteine	Dnmcys	N-(3,3-diphenylpropyl)glycine	Nbhe
	D-N-methylglutamine	Dnmgln	N-(3-guanidinopropyl)glycine	Narg
	D-N-methylglutamate	Dnmglu	N-(1-hydroxyethyl)glycine	Nthr
30	D-N-methylhistidine	Dnmhis	N-(hydroxyethyl)glycine	Nser

- 28 -

	D-N-methylisoleucine	Dnmile	N-(imidazolylethyl)glycine	Nhis
	D-N-methylleucine	Dnmleu	N-(3-indolylyethyl)glycine	Nhtrp
	D-N-methyllysine	Dnmlys	N-methyl- γ -aminobutyrate	Nmgabu
	N-methylcyclohexylalanine	Nmchexa	D-N-methylmethionine	Dnmmet
5	D-N-methylornithine	Dnmorn	N-methylcyclopentylalanine	Nmcpen
	N-methylglycine	Nala	D-N-methylphenylalanine	Dnmphe
	N-methylaminoisobutyrate	Nmaib	D-N-methylproline	Dnmpro
	N-(1-methylpropyl)glycine	Nile	D-N-methylserine	Dnmser
	N-(2-methylpropyl)glycine	Nleu	D-N-methylthreonine	Dnmthr
10	D-N-methyltryptophan	Dnmtrp	N-(1-methylethyl)glycine	Nval
	D-N-methyltyrosine	Dnmtyr	N-methyl- α -naphthylalanine	Nmanap
	D-N-methylvaline	Dnmval	N-methylpenicillamine	Nmpen
	γ -aminobutyric acid	Gabu	N-(<i>p</i> -hydroxyphenyl)glycine	Nhtyr
	L- <i>t</i> -butylglycine	Tbug	N-(thiomethyl)glycine	Ncys
15	L-ethylglycine	Etg	penicillamine	Pen
	L-homophenylalanine	Hphe	L- α -methylalanine	Mala
	L- α -methylarginine	Marg	L- α -methy lasparagine	Masn
	L- α -methy laspartate	Masp	L- α -methyl- <i>t</i> -butylglycine	Mtbug
	L- α -methylcysteine	Mcys	L-methylethylglycine	Metg
20	L- α -methylglutamine	Mgln	L- α -methylglutamate	Mglu
	L- α -methylhistidine	Mhis	L- α -methylhomophenylalanine	Mhphe
	L- α -methylisoleucine	Mile	N-(2-methylthioethyl)glycine	Nmet
	L- α -methylleucine	Mleu	L- α -methyllysine	Mlys
	L- α -methylmethionine	Mmet	L- α -methylnorleucine	Mnle
25	L- α -methylnorvaline	Mnva	L- α -methylornithine	Morn
	L- α -methylphenylalanine	Mphe	L- α -methylproline	Mpro
	L- α -methylserine	Mser	L- α -methylthreonine	Mthr
	L- α -methyltryptophan	Mtrp	L- α -methyltyrosine	Mtyr
	L- α -methylvaline	Mval	L-N-methylhomophenylalanine	Nmhph
30	N-(N-(2,2-diphenylethyl)	Nnbhm	N-(N-(3,3-diphenylpropyl)	Nnbhe

- 29 -

carbamylmethyl)glycine

carbamylmethyl)glycine

1-carboxy-1-(2,2-diphenyl- Nmbc
ethylamino)cyclopropane

-
- 5 Crosslinkers can be used, for example, to stabilise 3D conformations, using homo-bifunctional crosslinkers such as the bifunctional imido esters having $(CH_2)_n$ spacer groups with $n=1$ to $n=6$, glutaraldehyde, N-hydroxysuccinimide esters and hetero-bifunctional reagents which usually contain an amino-reactive moiety such as N-hydroxysuccinimide and another group specific-reactive moiety such as maleimido or
- 10 dithio moiety (SH) or carbodiimide (COOH). In addition, peptides can be conformationally constrained by, for example, incorporation of C_α and N_α -methylamino acids, introduction of double bonds between C_α and C_β atoms of amino acids and the formation of cyclic peptides or analogues by introducing covalent bonds such as forming an amide bond between the N and C termini, between two side chains or
- 15 between a side chain and the N or C terminus.

The nucleic acid molecule of the present invention is preferably in isolated form or ligated to a vector, such as an expression vector. By "isolated" is meant a nucleic acid molecule having undergone at least one purification step and this is conveniently defined,

20 for example, by a composition comprising at least about 10% subject nucleic acid molecule, preferably at least about 20%, more preferably at least about 30%, still more preferably at least about 40-50%, even still more preferably at least about 60-70%, yet even still more preferably 80-90% or greater of subject nucleic acid molecule relative to other components as determined by molecular weight, encoding activity, nucleotide

25 sequence, base composition or other convenient means. The nucleic acid molecule of the present invention may also be considered, in a preferred embodiment, to be biologically pure.

In a particularly preferred embodiment, the nucleotide sequence corresponding to *Bim* is

30 a cDNA sequence comprising a sequence of nucleotides as set forth in one of SEQ ID

- 30 -

NO: 1, 3 or 5 or is a derivative or homologue thereof including a nucleotide sequence having similarity to one of SEQ ID NO: 1, 3 or 5 and which encodes an amino acid sequence corresponding to an amino acid sequence as set forth in one of SEQ ID NO: 2, 4 or 6 or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

In another particularly preferred embodiment, the nucleotide sequence corresponding to *Bim* is a cDNA sequence comprising a sequence of nucleotides as set forth in one of SEQ ID NO: 7 or 9 or is a derivative or homologue thereof including a nucleotide sequence having similarity to one of SEQ ID NO: 7 or 9 and which encodes an amino acid sequence corresponding to an amino acid sequence as set forth in one of SEQ ID NO: 8 or 10 or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

15 A derivative of the nucleic acid molecule of the present invention also includes nucleic acid molecules capable of hybridising to the nucleotide sequences as set forth in one of SEQ ID NO: 1, 3, or 5 or SEQ ID NO: 7 or 9 under low stringency conditions.

Preferably, said low stringency is at 42°C.

20 In another embodiment the present invention is directed to an isolated nucleic acid molecule encoding *Bim* or a derivative thereof, said nucleic acid molecule selected from the list consisting of:

(i) A nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence set forth in one of SEQ ID NO: 2, 4, or 6 or a derivative or homologue thereof or having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.

(ii) A nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence set forth in one of SEQ ID NO: 8 or 10 or a derivative or

- 31 -

homologue or having at least about 45% similarity to one of SEQ ID NO: 8 or 10.

- (iii) A nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or 5 or a derivative or homologue thereof.
- (iv) A nucleic acid molecule comprising a nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9 or a derivative or homologue thereof.
- 10 (v) A nucleic acid molecule capable of hybridising under low stringency conditions at 42°C to the nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or 5 a derivative or homologue and encoding an amino acid sequence corresponding to an amino acid sequence as set forth in one of SEQ ID NO: 2, 4 or 6 a derivative or homologue or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- 15 (vi) A nucleic acid molecule capable of hybridising under low stringency conditions at 42°C to the nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9 a derivative or homologue and encoding an amino acid sequence corresponding to an amino acid sequence as set forth in one of SEQ ID NO: 8 or 10 a derivative or homologue or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.
- 20 (vii) A nucleic acid molecule capable of hybridising to the nucleic acid molecule of paragraphs (i) or (iii) or (v) under low stringency conditions at 42°C and encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- 25 (viii) A nucleic acid molecule capable of hybridising to the nucleic acid molecule of paragraphs (ii) or (iv) or (vi) under low stringency conditions at 42°C and
- 30

- 32 -

encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.

- (ix) A derivative or mammalian homologue of the nucleic acid molecule of paragraphs
5 (i) or (ii) or (iii) or (iv) or (v) or (vi) or (vii) or (viii).

The nucleic acid molecule may be ligated to an expression vector capable of expression in a prokaryotic cell (e.g. *E.coli*) or a eukaryotic cell (e.g. yeast cells, fungal cells, insect cells, mammalian cells or plant cells). The nucleic acid molecule may be ligated or fused
10 or otherwise associated with a nucleic acid molecule encoding another entity such as, for example, a signal peptide, a cytokine or other member of the Bcl-2 family.

The present invention extends to the expression product of the nucleic acid molecule hereinbefore defined.

15

The expression product is Bim having an amino acid sequence set forth in one of SEQ ID NO: 2, 4, 6, 8 or 10 or is a derivative or homologue thereof as defined above or is a mammalian homologue having an amino acid sequence of at least about 45% similarity to the amino acid sequence set forth in one of SEQ ID NO: 2, 4, 6, 8 or 10 or derivative or
20 homologue thereof.

Another aspect of the present invention is directed to an isolated polypeptide selected from the list consisting of:

- 25 (i) A polypeptide having an amino acid sequence substantially as set forth in one of SEQ ID NO: 2, 4, or 6 or derivative or homologue thereof or a sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- (ii) A polypeptide having an amino acid sequence substantially as set forth in one of
30 SEQ ID NO: 8 or 10 a derivative or homologue or a sequence having at least

- 33 -

about 45% similarity to one or more of SEQ ID NO: 8 or 10.

- (iii) A polypeptide encoded by a nucleotide sequence substantially as set forth in one of SEQ ID NO: 1, 3, or 5 or derivative or homologue thereof or a sequence
5 encoding an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- (iv) A polypeptide encoded by a nucleotide sequence substantially as set forth in one of SEQ ID NO: 7 or 9 or derivative or homologue thereof or a sequence encoding
10 an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 8 or 10.
- (v) A polypeptide encoded by a nucleic acid molecule capable of hybridising to the nucleotide sequence as set forth in one of SEQ ID NO: 1, 3, or 5 or derivative or
15 homologue thereof under low stringency conditions at 42°C and which encodes an amino acid sequence substantially as set forth in SEQ ID NO: 2, 4, or 6 or derivative or homologue thereof or an amino acid sequence having at least about 45% similarity to one or more of SEQ ID NO: 2, 4, or 6.
- 20 (vi) A polypeptide encoded by a nucleic acid molecule capable of hybridising to the nucleotide sequence as set forth in one of SEQ ID NO: 7 or 9 or derivative or homologue thereof under low stringency conditions at 42°C and which encodes an amino acid sequence substantially as set forth in SEQ ID NO: 8 or 10 or derivative or homologue thereof or an amino acid sequence having at least about
25 45% similarity to one or more of SEQ ID NO: 8 or 10.
- (vii) A polypeptide as defined in paragraphs (i) or (ii) or (iii) or (iv) or (v) or (vi) in homodimeric form.
- 30 (viii) A polypeptide as defined in paragraphs (i) or (ii) or (iii) or (iv) or (v) or (vi) in

- 34 -

heterodimeric form.

As defined earlier, the present invention extends to peptides or derivatives thereof of Bim. Preferably, said peptide comprises at least 5 contiguous amino acids of the
5 polypeptide defined in SEQ ID NO:2, 4, 6, 8 or 10. The present invention also extends to nucleic acid molecules encoding the peptides of the present invention.

Another aspect of the present invention provides a nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide having one or more of the identifying
10 characteristics of Bim or a derivative or homologue thereof.

Reference herein to "identifying characteristics" of Bim includes one or more of the following features:

- 15 (i) A polypeptide which induces apoptosis.
- (ii) A polypeptide having an amino acid sequence substantially as set forth in SEQ ID NO:2, 4, 6, 8 or 10 or a derivative or homologue thereof.
- 20 (iii) A polypeptide having an amino acid sequence of at least 45% similarity to SEQ ID NO:2, 4, 6, 8 or 10.
- (iv) A polypeptide as defined in paragraph (ii) or (iii) which induces apoptosis.
- 25 (v) A polypeptide encoded by a nucleic acid sequence substantially as set forth in SEQ ID NO:1, 3, 5, 7 or 9 or derivative or homologue thereof.
- (vi) A polypeptide encoded by a nucleic acid molecule capable of hybridising to the nucleotide sequence as set forth in one of SEQ ID NO: 1, 3, 5, 7 or 9 under low
30 stringency conditions at 42°C.

- 35 -

- (vii) A polypeptide as defined in paragraph (v) or (vi) which induces apoptosis.
- (viii) A non-apoptosis inducing derivative of the polypeptide defined in paragraphs (i) to (vii).

5

The present invention should be understood to extend to the expression product of the nucleic acid molecule according to this aspect of the present invention.

Although not intending to limit the invention to any one theory or mode of action, the BH3 region is responsible for some of the cytotoxic actions of Bim. The BH3 region forms an amphipathic helix that interacts with the elongated hydrophobic cleft formed by the BH1, BH2 and BH3 regions of pro-survival molecules such as, for example, Bcl-x_L. The pro-apoptotic action of Bim reflects its ability to bind to the anti-apoptotic members of the Bcl-2 family. Bim is the only BH3-only protein for which splice variants have been described. Isoforms such as Bim_S, Bim_L and Bim_{EL} interact *in vivo* with Bcl-2 family members but induce cell death with different degrees of cytotoxicity. Bim_S, for example, is a more potent inducer of cell death than Bim_L or Bim_{EL}.

Still without limiting the invention to any one theory or mode of action, the pro-apoptotic activity of Bim is thought to be regulated both at the transcriptional level and at the post-translational level. Sequence analysis of the non-coding 5' region of *Bim* has revealed a number of putative binding sites for transcription factors. Bim_L and Bim_{EL} can bind to dynein light chain. Dynein light chain is a highly conserved protein which is a component of the dynein motor complex. The dynein motor complex moves vesicles along microtubules but may also carry out other functions.

In living cells Bim is bound to the dynein motor complex and associated with the microtubular network. When cells are stressed, for example by removal of growth factors or UV irradiation, Bim is rapidly released from the dynein motor complex, but remains still bound to dynein light chain. Thus the breakage occurs between dynein light

- 36 -

chain and dynein intermediate chain. This change in subcellular localisation is thought to constitute an upstream signalling event, probably for cell death.

The interaction of Bim with the dynein motor complex regulates the pro-apoptotic activity of Bim. It is thought that when Bim is released from the microtubular network it is free to interact with Bcl-2 and its homologues and will thereby prevent their pro-survival function. Consistent with this idea, Bim_S, which does not bind to dynein light chain, is not associated with the microtubular network and is a much more potent killer than Bim_L or Bim_{EL}. Single amino acid mutations in Bim that abolish binding to dynein light chain have been identified.

Accordingly, a related aspect of the present invention is directed to a variant of an isolated *Bim* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion to the polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

Preferably, the mutation results in an altered amino acid sequence in the region which binds the dynein light chain. For example, in murine and human Bim_L this corresponds to the region defined by amino acid residue numbers 42 to 71, in murine Bim_{EL} this region is defined by amino acid residue numbers 42 to 127 and in human Bim_{EL} amino acid residue numbers 42 to 131. The present invention should be understood to extend to variants of Bim comprising a mutation resulting in an amino acid addition, substitution and/or deletion in a region functionally equivalent to the regions hereinbefore defined.

Accordingly, the present invention is more particularly directed to a variant of an isolated *Bim* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion in the region of the polypeptide encoded by said variant which binds the dynein light chain wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light

- 37 -

chain.

Even more preferably, the present invention is directed to a variant of an isolated murine or human *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations in
 5 said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion in the region defined by amino acid residue numbers 42 to 71 of the polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

- 10 In another preferred embodiment the present invention is directed to a variant of an isolated murine *Bim_{EL}* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution and/or deletion in the region defined by amino acid residue numbers 42 to 127 of the polypeptide encoded by said variant wherein said polypeptide cannot bind,
 15 couple or otherwise associate with a dynein light chain.

In yet another preferred embodiment the present invention is directed to a variant of an isolated human *Bim_{EL}* nucleic acid molecule comprising one or more nucleotide mutations in said nucleic acid molecule resulting in at least one amino acid addition, substitution
 20 and/or deletion in the region defined by amino acid residue numbers 42 to 131 of the polypeptide encoded by said variant wherein said polypeptide cannot bind, couple or otherwise associate with a dynein light chain.

Mutations contemplated by the present invention which occur in combination with one or
 25 more mutations in another location are also contemplated by the present invention.

Preferably, the nucleotide mutation is a mutation to the human or murine *Bim_L* and results in an amino acid substitution of one or more of D51, S53, T54 and/or N65. Preferred mutations include one or more of D51G, S53P, T54A, T54I and/or N65S.

Accordingly, a preferred embodiment of the present invention is directed to a variant of an isolated human or murine *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitution D51G of the polypeptide encoded by said mutated nucleic acid molecule.

5

Another preferred embodiment of the present invention is directed to a variant of an isolated human or murine *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitution S53P of the polypeptide encoded by said mutated nucleic acid molecule.

10

In another preferred embodiment the present invention provides a variant of an isolated human or murine *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitution T54A of the polypeptide encoded by said mutated nucleic acid molecule.

15

In yet another preferred embodiment the present invention provides a variant of an isolated *Bim_L* nucleic acid molecule comprising one or more nucleotide mutations resulting in the amino acid substitutions T54I and N65S of the polypeptide encoded by said mutated nucleic acid molecule.

20

The present invention extends to the expression products of the nucleic acid molecule variants defined according to this aspect of the present invention.

Accordingly, the present invention is directed to a variant of an isolated Bim polypeptide comprising at least one amino acid addition, substitution and/or deletion wherein said polypeptide cannot bind, couple or otherwise associate with the dynein light chain.

Preferably said addition, substitution and/or deletion is of any one or more amino acid residues located in the region which binds the dynein light chain. Even more preferably said region is defined by residue numbers 42 to 71 in murine and human *Bim_L*, residue

30

- 39 -

numbers 42 to 127 in murine Bim_{EL} and residue numbers 42 to 131 in human Bim_{EL}.

Preferably said amino acid addition, substitution and/or deletion is a substitution of D51, S53, T54 and/or N65 of human or murine Bim_L. Preferred mutations include one or
5 more of D51G, S53P, T54A, T54I and N65S. Most preferably said mutation is D51G or S53P or T54A or T54I and N65S.

The present invention extends to derivatives of the nucleic acid molecules and polypeptides according to this aspect of the present invention. The term "derivatives"
10 should be understood as previously defined.

As hereinbefore defined, reference to "Bim" and "*Bim*" should be understood to include reference to the variant molecules defined according to this aspect of the present
15 invention.

The Bim of the present invention may be in multimeric form meaning that two or more molecules are associated together. Where the same Bim molecules are associated together, the complex is a homomultimer. An example of a homomultimer is a
20 homodimer. Where at least one Bim is associated with at least one non-Bim molecule, then the complex is a heteromultimer such as a heterodimer. A heteromultimer may include a molecule of another member of the Bcl-2 family or other molecule capable of modulating apoptosis.

25 The present invention contemplates, therefore, a method for modulating expression of *Bim* in a mammal, said method comprising administering to said mammal a modulating effective amount of an agent for a time and under conditions sufficient to up-regulate or down-regulate or otherwise modulate expression of *Bim*. For example, *Bim* antisense sequences such as oligonucleotides may be introduced into a cell to enhance the ability of
30 that cell to survive. Conversely, a nucleic acid molecule encoding Bim or a derivative

- 40 -

thereof may be introduced to decrease the survival capacity of any cell expressing the endogenous *Bim* gene. Modulation of the expression of *Bim* should be understood to extend to modulating transcriptional and translation events such as the splicing pattern of *Bim* RNA.

5

Another aspect of the present invention contemplates a method of modulating activity of *Bim* in a mammal, said method comprising administering to said mammal a modulating effective amount of an agent for a time and under conditions sufficient to increase or decrease *Bim* activity.

10

Modulation of said activity by the administration of an agent to a mammal can be achieved by one of several techniques, including but in no way limited to introducing into said mammal a proteinaceous or non-proteinaceous molecule which:

15

- (i) modulates expression of *Bim*;
- (ii) functions as an antagonist of *Bim*;
- (iii) functions as an agonist of *Bim*.

20

Said proteinaceous molecule may be derived from natural or recombinant sources including fusion proteins or following, for example, natural product screening. Said non-proteinaceous molecule may be, for example, a nucleic acid molecule or may be derived from natural sources, such as for example natural product screening or may be
25 chemically synthesised. The present invention contemplates chemical analogues of *Bim* capable of acting as agonists or antagonists of *Bim*. Chemical agonists may not necessarily be derived from *Bim* but may share certain conformational similarities. Alternatively, chemical agonists may be specifically designed to mimic certain physiochemical properties of *Bim*. Antagonists may be any compound capable of
30 blocking, inhibiting or otherwise preventing *Bim* from carrying out its normal or

- 41 -

pathological biological functions. Antagonists include, but are not limited to parts of Bim or peptides thereof, monoclonal antibodies specific for Bim or parts of Bim, and antisense nucleic acids or oligonucleotides which prevent transcription or translation of *Bim* genes or mRNA in mammalian cells. Agonists of Bim and *Bim* include, for example, the derivative or variant molecules or peptides hereinbefore defined which interact with anti-apoptotic molecules such as Bcl-2, to prevent their functional activity thereby promoting apoptosis. Agonists may also include molecules capable of disrupting or preventing binding of Bim to the dynein light chain or the interaction of dynein light chain with dynein intermediate chain.

10

Said proteinaceous or non-proteinaceous molecule may act either directly or indirectly to modulate the expression of *Bim* or the activity of Bim. Said molecule acts directly if it associates with *Bim* or Bim to modulate the expression or activity of *Bim* or Bim. Said molecule acts indirectly if it associates with a molecule other than *Bim* or Bim which other molecule either directly or indirectly modulates the expression or activity of *Bim* or Bim. Accordingly, the method of the present invention encompasses the regulation of *Bim* or Bim expression or activity via the induction of a cascade of regulatory steps which lead to the regulation of *Bim* or Bim expression or activity.

Increased *Bim* expression or Bim activity is useful, for example, for treatment or prophylaxis in conditions such as cancer and deletion of autoreactive lymphocytes in autoimmune disease. Decreased *Bim* expression or Bim activity is useful in regulating inhibition or prevention of cell death or degeneration such as under cytotoxic conditions during, for example, γ -irradiation and chemotherapy or during HIV/AIDS or other viral infections, ischaemia or myocardial infarction. Since Bim is expressed in germ cells, modulating *Bim* expression or Bim activity is useful, for example, as a contraceptive or method of sterilisation by preventing generation of fertile sperm.

Another aspect of the present invention contemplates a method of modulating apoptosis in a mammal said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate the expression of a

30

- 42 -

nucleotide sequence encoding *Bim*.

Yet another aspect of the present invention contemplates a method of modulating apoptosis in a mammal said method comprising administering to said mammal an
5 effective amount of an agent for a time and under conditions sufficient to modulate the activity of *Bim*.

Still another aspect of the present invention contemplates a method of modulating apoptosis in a mammal said method comprising administering to said mammal an
10 effective amount of *Bim* or *Bim* or derivative thereof.

The *Bim*, *Bim* or derivative thereof or agent used may also be linked to a targeting means such as a monoclonal antibody, which provides specific delivery of the *Bim*, *Bim* or agent to the target cells.

15

In a preferred embodiment of the present invention, the *Bim*, *Bim* or agent used in the method is linked to an antibody specific for said target cells to enable specific delivery to these cells.

20 Administration of the *Bim*, *Bim* or agent, in the form of a pharmaceutical composition, may be performed by any convenient means. *Bim*, *Bim* or agent of the pharmaceutical composition are contemplated to exhibit therapeutic activity when administered in an amount which depends on the particular case. The variation depends, for example, on the human or animal and the *Bim*, *Bim* or agent chosen. A broad range of doses may be
25 applicable. Considering a patient, for example, from about 0.01 mg to about 10 mg of *Bim* or agent may be administered per kilogram of body weight per day. Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily, weekly, monthly or other suitable time intervals or the dose may be proportionally reduced as indicated by the exigencies of the
30 situation. The *Bim* or agent may be administered in a convenient manner such as by the

- 43 -

oral, intravenous (where water soluble), intranasal, intraperitoneal, intramuscular, subcutaneous, intradermal or suppository routes or implanting (e.g. using slow release molecules). With particular reference to use of Bim or agent, these peptides may be administered in the form of pharmaceutically acceptable nontoxic salts, such as acid addition salts or metal complexes, e.g. with zinc, iron or the like (which are considered as salts for purposes of this application). Illustrative of such acid addition salts are hydrochloride, hydrobromide, sulphate, phosphate, maleate, acetate, citrate, benzoate, succinate, malate, ascorbate, tartrate and the like. If the active ingredient is to be administered in tablet form, the tablet may contain a binder such as tragacanth, corn starch or gelatin; a disintegrating agent, such as alginic acid; and a lubricant, such as magnesium stearate.

A further aspect of the present invention relates to the use of the invention in relation to mammalian disease conditions. For example, the present invention is particularly applicable to, but in no way limited to, use in therapy or prophylaxis in relation to cancer, degenerative diseases, autoimmune disorders, viral infections or for germ cell regulation.

Accordingly, another aspect of the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate the expression of *Bim* wherein said modulation results in modulation of apoptosis.

In another aspect the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate the activity of Bim wherein said modulation results in modulation of apoptosis.

In another aspect the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of Bim or

- 44 -

derivative thereof for a time and under conditions sufficient to modulate apoptosis.

Yet another aspect the present invention relates to a method of treating a mammal said method comprising administering to said mammal an effective amount of *Bim* or
5 derivative thereof for a time and under conditions sufficient to modulate apoptosis.

In yet another aspect the present invention relates to the use of an agent capable of modulating the expression of *Bim* or derivative thereof in the manufacture of a medicament for the modulation of apoptosis.

10

Another aspect of the present invention relates to the use of an agent capable of modulating the expression of *Bim* or derivative thereof in the manufacture of a medicament for the modulation of apoptosis.

15 A further aspect of the present invention relates to the use of *Bim* or *Bim* or derivative thereof in the manufacture of a medicament for the modulation of apoptosis.

Still yet another aspect of the present invention relates to agents for use in modulating *Bim* expression wherein modulating expression of said *Bim* modulates apoptosis.

20

A further aspect of the present invention relates to agents for use in modulating *Bim* expression wherein modulating expression of said *Bim* modulates apoptosis.

Another aspect of the present invention relates to *Bim* or *Bim* or derivative thereof for
25 use in modulating apoptosis.

In a related aspect of the present invention, the mammal undergoing treatment may be human or an animal in need of therapeutic or prophylactic treatment.

30 In yet another further aspect the present invention contemplates a pharmaceutical

composition comprising *Bim*, Bim or derivative thereof or an agent capable of modulating *Bim* expression or Bim activity together with one or more pharmaceutically acceptable carriers and/or diluents. *Bim*, Bim or said agent are referred to as the active ingredients.

5

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water soluble) and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of
10 manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a
15 coating such as licithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The preventions of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or
20 sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the
25 required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable
30 solutions, the preferred methods of preparation are vacuum drying and the freeze-drying

technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

When *Bim*, Bim and Bim modulators are suitably protected they may be orally
5 administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsule, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups,
10 wafers, and the like. Such compositions and preparations should contain at least 1 % by weight of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically useful compositions in such that a suitable dosage will be obtained. Preferred compositions or
15 preparations according to the present invention are prepared so that an oral dosage unit form contains between about 0.1 μ g and 2000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the following: A binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as
20 dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other
25 materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should
30 be pharmaceutically pure and substantially non-toxic in the amounts employed. In

- 47 -

addition, the active compound may be incorporated into sustained-release preparations and formulations.

Pharmaceutically acceptable carriers and/or diluents include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active material for the treatment of disease in living subjects having a diseased condition in which bodily health is impaired as herein disclosed in detail.

The principal active ingredient is compounded for convenient and effective administration in effective amounts with a suitable pharmaceutically acceptable carrier in dosage unit form as hereinbefore disclosed. A unit dosage form can, for example, contain the principal active compound in amounts ranging from 0.5 μ g to about 2000 mg. Expressed in proportions, the active compound is generally present in from about 0.5 μ g to about 2000 mg/ml of carrier. In the case of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of

- 48 -

administration of the said ingredients.

The pharmaceutical composition may also comprise genetic molecules such as a vector capable of transfecting target cells where the vector carries a nucleic acid molecule
5 capable of modulating *Bim* expression or Bim activity. The vector may, for example, be a viral vector.

Conditions requiring modulation of physiological cell death include enhancing survival of cells utilising, for example, antisense sequence in patients with neurodegenerative
10 diseases, myocardial infarction, muscular degenerative disease, hypoxia, ischaemia, HIV infection or for prolonging the survival of cells being transplanted for treatment of disease. Alternatively, the molecules of the present invention are useful for, for example, reducing the survival capacity of tumour cells or autoreactive lymphocytes.

The anti-sense sequence may also be used for modifying *in vitro* behaviour of cells, for
15 example, as part of a protocol to develop novel lines from cell types having unidentified growth factor requirements; for facilitating isolation of hybridoma cells producing monoclonal antibodies, as described below; and for enhancing survival of cells from primary explants while they are being genetically modified.

20 Still another aspect of the present invention is directed to an immunointeractive molecule comprising an antigen binding portion having specificity for Bim or *Bim* or derivative thereof.

Reference to "immunointeractive molecule" should be understood as a reference to any
25 molecule comprising an antigen binding portion or a derivative of said molecule.

Examples of molecules contemplated by this aspect of the present invention include, but are not limited to, monoclonal and polyclonal antibodies (including synthetic antibodies, hybrid antibodies, humanized antibodies, catalytic antibodies) and T cell antigen binding molecules. Preferably, said immunoreactive molecule is a monoclonal antibody.

- 49 -

According to this preferred embodiment there is provided a monoclonal antibody having specificity for Bim or *Bim* or derivative thereof.

Reference to a molecule "having specificity for Bim or *Bim*" should be understood as a
5 reference to a molecule, such as a monoclonal antibody, having specificity for any one or more epitopes of Bim or *Bim*. These epitopes may be conformational epitopes, linear epitopes or a combination of conformational and linear epitopes of either the native Bim or *Bim* molecule or the denatured molecule.

10 More preferably there is provided a monoclonal antibody having specificity for Bim_L.

The immunointeractive molecules of the present invention may be naturally occurring, synthetic or recombinantly produced. For example, monoclonal or polyclonal antibodies may be selected from naturally occurring antibodies to Bim or *Bim* or may be specifically
15 raised to Bim or *Bim*. In the case of the latter, Bim or *Bim* may first need to be associated with a carrier molecule. The antibodies and/or recombinant Bim of the present invention are particularly useful as therapeutic or diagnostic agents.

Alternatively, fragments of antibodies may be used such as Fab fragments. Furthermore, the present invention extends to recombinant and synthetic antibodies, to antibody
20 hybrids and to antibodies raised against non-Bim antigens but which are cross-reactive with any one or more Bim epitopes. A "synthetic antibody" is considered herein to include fragments and hybrids of antibodies. The antibodies of this aspect of the present invention are particularly useful for immunotherapy and may also be used as a diagnostic tool for assessing apoptosis or monitoring the program of a therapeutic regime.

25

For example, Bim and *Bim* can be used to screen for naturally occurring antibodies to Bim and *Bim*, respectively. These may occur, for example in some degenerative disorders.

30 For example, specific antibodies can be used to screen for Bim proteins. The latter

- 50 -

would be important, for example, as a means for screening for levels of Bim in a cell extract or other biological fluid or purifying Bim made by recombinant means from culture supernatant fluid. Techniques for the assays contemplated herein are known in the art and include, for example, sandwich assays, ELISA and flow cytometry.

5

It is within the scope of this invention to include any second antibodies (monoclonal, polyclonal or fragments of antibodies) directed to the first mentioned antibodies discussed above. Both the first and second antibodies may be used in detection assays or a first antibody may be used with a commercially available anti-immunoglobulin antibody. An antibody as contemplated herein includes any antibody specific to any region of Bim.

Both polyclonal and monoclonal antibodies are obtainable by immunization with the protein or peptide derivatives and either type is utilizable for immunoassays. The methods of obtaining both types of sera are well known in the art. Polyclonal sera are less preferred but are relatively easily prepared by injection of a suitable laboratory animal with an effective amount of Bim, or antigenic parts thereof, collecting serum from the animal, and isolating specific sera by any of the known immunoabsorbent techniques. Although antibodies produced by this method are utilizable in virtually any type of immunoassay, they are generally less favoured because of the potential heterogeneity of the product.

The use of monoclonal antibodies in an immunoassay is particularly preferred because of the ability to produce them in large quantities and the homogeneity of the product. The preparation of hybridoma cell lines for monoclonal antibody production derived by fusing an immortal cell line and lymphocytes sensitized against the immunogenic preparation can be done by techniques which are well known to those who are skilled in the art. (See, for example Douillard and Hoffman, Basic Facts about Hybridomas, in *Compendium of Immunology* Vol II, ed. by Schwartz, 1981; Kohler and Milstein, *Nature* 256: 495-499, 1975; *European Journal of Immunology* 6: 511-519, 1976).

30

- 51 -

Screening for immunointeractive molecules, such as antibodies, can be a time consuming and labour intensive process. However, the inventors have developed a rapid and efficient flow cytometric screening procedure for the identification of immunointeractive molecules, and in particular antibodies, directed to low abundance cytoplasmic proteins
5 such as, but not limited to, Bim.

The method according to this aspect of the present invention is based on the analysis of a population of cells, following the incubation of these cells with the antibody of interest together with or separately to a reporter molecule, said population of cells comprising
10 both cells expressing the protein of interest and cells which do not express the protein of interest. This analysis is preferably flow cytometric analysis and the cells expressing the protein of interest are preferably transfected with a nucleic acid molecule encoding the protein of interest to thereby express high levels of said protein. Where the protein is a cytoplasmic protein the cells are permeabilised prior to incubation with the antibody of
15 interest. By screening a population of cells comprising both cells which do not express and cells which do express the protein of interest, determination of which antibodies bind to the protein of interest is simplified since where the subject antibody is directed to the protein of interest, a double fluorescence peak is observed. The lower intensity peak represents background staining while the higher fluorescence intensity peak is the result of
20 specific staining. Where the antibody being screened according to this method is not directed to the protein of interest, a single peak of low fluorescence intensity is observed. Antibodies not specific to the protein of interest but bound to some unknown epitope present in both populations of cells produces a single peak with high fluorescence intensity. This technique provides a rapid and accurate method of screening for
25 immunointeractive molecules directed to low abundance intracytoplasmic molecules.

Accordingly, another aspect of the present invention provides a method of detecting an immunointeractive molecule, in a sample, specific for a protein of interest produced by a cell said method comprising contacting the sample to be tested with a population of cells
30 comprising a defined ratio of cells producing the protein of interest and cells not

- 52 -

producing the protein of interest for a time and under conditions sufficient for immunointeractive molecules, if present in said sample, to interact with said protein of interest and the subjecting said immunointeractive molecule-protein complex to detecting means.

5

Preferably said immunointeractive molecule is an antibody.

More preferably, said detecting means comprises an antiimmunoglobulin antibody labelled with a reporter molecule capable of giving a detectable signal. Even more preferably
10 said reporter molecule is fluorochrome.

Reference to "sample" should be understood as a reference to any sample potentially comprising an immunointeractive molecule, such as an antibody. Said immunointeractive molecule may be produced by natural, recombinant or synthetic
15 means.

The method of the present invention is predicated on subjecting the cells incubated with the sample of the present invention to flow cytometric analysis to produce a fluorescent signal wherein a differential fluorescent signal is indicative of antibody binding to the
20 target protein expressed by said cells.

The method exemplified herein is directed, but not limited to, screening for immunointeractive molecules comprising an antigen binding site directed to epitopes of Bim. The promyelomonocytic cell line FDC-P1 is transfected with a Bcl-2 expression
25 construct and an EE (Glu-Glu) epitope-tagged Bim construct. A 1:1 ratio of Bcl-2 transfected cells to Bim transfected cells are fixed, permeabilised and contacted with the immunointeractive molecule of interest, such as a hybridoma supernatant. Visualisation of antibodies bound intracellular molecules can be achieved via a number of techniques known to those skilled in the art, including, for example, the use of fluorescently labelled
30 reporter molecules. Where the antibody of interest is directed to Bim, a double

- 53 -

fluorescence peak is observed, the lower intensity peak representing background staining of the Bcl-2 transfected negative control cells.

In another aspect of the present invention, the molecules of the present invention are also
5 useful as screening targets for use in applications such as the diagnosis of disorders which
are regulated by Bim. For example, screening for the levels of Bim or *Bim* in tissue as
an indicator of a predisposition to, or the development or, cancer, a degenerative disease
or infertility. The screening of this aspect of the present invention may also be directed
to detecting mutations in Bim or *Bim*.

10

Accordingly, another aspect of the present invention contemplates a method for detecting
Bim in a biological sample from a subject said method comprising contacting said
biological sample with an immunointeractive molecule as hereinbefore defined specific
for Bim or its derivatives thereof for a time and under conditions sufficient for an
15 immunointeractive molecule-Bim complex to form, and then detecting said complex.

Preferably said immunointeractive molecule is an antibody. Even more preferably said
antibody is a monoclonal antibody.

20 Reference to biological sample according to this aspect of the present invention should be
understood as a reference to any sample comprising tissue from a subject said "tissue"
should be understood in its broadest sense to include biological fluid, biopsy samples or
any other form of tissue or fluid or extracts therefrom such as DNA or RNA properties.

25 Still another aspect of the present invention contemplates a method for detecting *Bim* in a
biological sample from a subject said method comprising contacting said biological
sample with an immunointeractive molecule as hereinbefore defined specific for *Bim* or
its derivatives thereof for a time and under conditions sufficient for an immunointeractive
molecule-*Bim* complex to form, and then detecting said complex.

30

- 54 -

Reference to an "immunointeractive" molecule should be understood as a reference to any molecule which couples, binds or otherwise associates with *Bim* or Bim or derivative thereof. For example said interactive molecule may be a nucleic acid molecule or an anti-nuclear antibody.

5

The presence of Bim may be determined in a number of ways such as by Western blotting, ELISA or flow cytometry procedures. *Bim* mRNA or DNA may be detected, for example, by *in situ* hybridization or Northern blotting or Southern blotting. These, of course, include both single-site and two-site or "sandwich" assays of the non-competitive types, as well as in the traditional competitive binding assays. These assays also include direct binding of a labelled antibody to a target.

Sandwich assays are among the most useful and commonly used assays and are favoured for use in the present invention. A number of variations of the sandwich assay technique exist, and all are intended to be encompassed by the present invention. Briefly, in a typical forward assay, an unlabelled antibody is immobilized on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen complex, a second antibody specific to the antigen, labelled with a reporter molecule capable of producing a detectable signal is then added and incubated, allowing time sufficient for the formation of another complex of antibody-antigen-labelled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal produced by the reporter molecule. The results may either be qualitative, by simple observation of the visible signal, or may be quantitated by comparing with a control sample containing known amounts of hapten. Variations on the forward assay include a simultaneous assay, in which both sample and labelled antibody are added simultaneously to the bound antibody. These techniques are well known to those skilled in the art, including any minor variations as will be readily apparent. In accordance with the present invention the sample is one which might contain Bim including cell extract, tissue biopsy or possibly serum, saliva, mucosal secretions, lymph,

- 55 -

tissue fluid and respiratory fluid. The sample is, therefore, generally a biological sample comprising biological fluid but also extends to fermentation fluid and supernatant fluid such as from a cell culture.

5 In the typical forward sandwich assay, a first antibody having specificity for the Bim or antigenic parts thereof, is either covalently or passively bound to a solid surface. The solid surface is typically glass or a polymer, the most commonly used polymers being cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene. The solid supports may be in the form of tubes, beads, discs of microplates, or any other
10 surface suitable for conducting an immunoassay. The binding processes are well-known in the art and generally consist of cross-linking covalently binding or physically adsorbing, the polymer-antibody complex is washed in preparation for the test sample. An aliquot of the sample to be tested is then added to the solid phase complex and incubated for a period of time sufficient (e.g. 2-40 minutes) and under suitable conditions
15 (e.g. 25°C) to allow binding of any subunit present in the antibody. Following the incubation period, the antibody subunit solid phase is washed and dried and incubated with a second antibody specific for a portion of the hapten. The second antibody is linked to a reporter molecule which is used to indicate the binding of the second antibody to the hapten.

20

An alternative method involves immobilizing the target molecules in the biological sample and then exposing the immobilized target to specific antibody which may or may not be labelled with a reporter molecule. Depending on the amount of target and the strength of the reporter molecule signal, a bound target may be detectable by direct
25 labelling with the antibody. Alternatively, a second labelled antibody, specific to the first antibody is exposed to the target-first antibody complex to form a target-first antibody-second antibody tertiary complex. The complex is detected by the signal emitted by the reporter molecule.

30 By "reporter molecule" as used in the present specification, is meant a molecule which,

- 56 -

by its chemical nature, provides an analytically identifiable signal which allows the detection of antigen-bound antibody. Detection may be either qualitative or quantitative. The most commonly used reporter molecules in this type of assay are either enzymes, fluorophores or radionuclide containing molecules (i.e. radioisotopes) and
5 chemiluminescent molecules.

In the case of an enzyme immunoassay, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. As will be readily recognized, however, a wide variety of different conjugation techniques exist, which are readily
10 available to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase and alkaline phosphatase, amongst others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable color change. Examples of suitable enzymes include alkaline phosphatase and peroxidase. It is also possible to
15 employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. In all cases, the enzyme-labelled antibody is added to the first antibody hapten complex, allowed to bind, and then the excess reagent is washed away. A solution containing the appropriate substrate is then added to the complex of antibody-antigen-antibody. The substrate will react with the enzyme linked
20 to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of hapten which was present in the sample. "Reporter molecule" also extends to use of cell agglutination or inhibition of agglutination such as red blood cells on latex beads, and the like.

25

Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labelled antibody adsorbs the light energy, inducing a state of excitability in the molecule, followed by
30 emission of the light at a characteristic color visually detectable with a light microscope.

- 57 -

As in the EIA, the fluorescent labelled antibody is allowed to bind to the first antibody-hapten complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to the light of the appropriate wavelength the fluorescence observed indicates the presence of the hapten of interest. Immunofluorescence and EIA techniques
5 are both very well established in the art and are particularly preferred for the present method. However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules, may also be employed.

The present invention also contemplates genetic assays such as involving PCR analysis to
10 detect *Bim* or its derivatives.

Further features of the present invention are more fully described in the following examples. It is to be understood, however, that this detailed description is included solely for the purposes of exemplifying the present invention. It should not be
15 understood in any way as a restriction on the broad description of the invention as set out above.

SUMMARY OF SEQ ID NO:

Sequence	SEQ ID NO:
5 nucleotide sequence of murine Bim _S	1 ←
amino acid sequence of murine Bim _S	2
nucleotide sequence of murine Bim _L	3 ←
amino acid sequence of murine Bim _L	4
nucleotide sequence of murine Bim _{EL}	5 ←
10 amino acid sequence of murine Bim _{EL}	6
nucleotide sequence of human Bim _L	7 ←
amino acid sequence of human Bim _L	8
nucleotide sequence of human Bim _{EL}	9 ←
amino acid sequence of human Bim _{EL}	10
15 peptides	11-13
oligonucleotide primers	14-26

EXAMPLE 1

Isolation of a novel gene encoding a Bcl-2-binding protein

In an attempt to identify novel proteins that bind to Bcl-2, we used recombinant human
5 Bcl-2 protein, labelled with ^{32}P (Blancar and Rutter, 1992), to screen a bacteriophage λ
cDNA expression library constructed from the p53^{-/-} T lymphoma cell line KO52DA20
(Strasser *et al.*, 1994). A screen of 10^6 clones yielded 5 independent clones which
encoded the same novel protein, which we named Bim, for Bcl-2 interacting mediator of
cell death. Sequence analysis of the *bim* cDNAs revealed three variants of the coding
10 region, apparently produced by alternative splicing (Figure 1A). Reverse transcriptase-
PCR on mRNA from KO52DA20 cells gave PCR products of the sizes expected for each
of these transcripts, which we designated *bim*_{EL}, *bim*_L and *bim*_S, although the last was in
low yield (data not shown). The predicted proteins Bim_{EL}, Bim_L and Bim_S comprise 196,
140 and 110 amino acid residues (Figure 1B). Hybridising human embryo and liver
15 cDNA libraries with mouse *bim* cDNA yielded human cDNAs encoding Bim_L and Bim_{EL}.
Human Bim_{EL} is a protein of 198 residues, 89% identical to its mouse counterpart
(Figure 1C), and human Bim_L (138 residues) is 85% identical to mouse Bim_L.

Bim has no substantial homology with any protein in current databases. However,
20 scrutiny of its sequence (Figure 1C) revealed a stretch of nine amino acids corresponding
to a BH3 homology region (Boyd *et al.*, 1995; Chittenden *et al.*, 1995). Apart from this
region, the Bim sequence is unrelated to that of any other BH3-containing protein; it
contains no other BH region, nor indeed any other known functional motif. The protein
does have a C-terminal hydrophobic region (Figure 1C), raising the possibility that it
25 associates with membranes.

Northern blot analysis showed that *bim* was expressed in a number of B and T lymphoid
cell lines, although not in the myeloid line FDC-P1 (Figure 2). A major transcript of 5.7
kb and minor transcripts of 3.8, 3.0, and 1.4 kb were detected. Neither the level nor
30 relative abundance of these transcripts changed significantly in KO52DA20 cells induced

- 60 -

to undergo apoptosis by treatment with dexamethasone (Figure 2, compare lanes 1 and 2, and lanes 3 and 4) or exposure to γ -radiation (compare lanes 1 and 5). Overexpression of *bcl-2* in several of the lines did not affect *bim* mRNA levels (Figure 2).

5

EXAMPLE 2

Bim localises to cytoplasmic membranes

The presence of the C-terminal hydrophobic domain in Bim prompted us to investigate its subcellular localisation. L929 fibroblasts were transiently transfected with an expression
10 vector encoding Bim_L tagged with an N-terminal EE-epitope, and the permeabilised cells were stained with an anti-EE monoclonal antibody. Confocal microscopy revealed that Bim_L was cytoplasmic and apparently associated with intracellular membranes (Figure 3A). We also introduced the *bim_L* vector into L929 cells stably infected with a human Bcl-2 encoding retrovirus (Lithgow *et al.*, 1994). The similarity of the anti-EE staining
15 pattern of these cells (Figure 3C) to that of those expressing Bim_L alone (Figure 3A) demonstrated that high concentrations of Bcl-2 did not perturb the localisation of Bim_L. The pattern of Bim_L staining was similar to that reported for Bcl-2 (Monaghan *et al.*, 1992; Krajewski *et al.*, 1993; Lithgow *et al.*, 1994), and overlaying the images obtained from the same cells stained with anti-Bcl-2 (Figure 3B) and anti-EE (Figure 3C)
20 antibodies showed that the two proteins co-localised (Figure 3D).

EXAMPLE 3

Overexpression of Bim kills cells by a pathway requiring caspases

25

Other known 'BH3-only' proteins (Bik/Nbk, Bid and Hrk) provoke apoptosis when highly expressed (Boyd *et al.*, 1995; Han *et al.*, 1996; Wang *et al.*, 1996; Inohara *et al.*, 1997). We therefore tested whether Bim is cytotoxic by transiently transfecting 293T human embryonal kidney cells with a plasmid encoding EE-Bim_L. The viability of the
30 transfected cells was determined subsequently by flow cytometric analysis of

- 61 -

permeabilised cells stained with the anti-EE antibody and the DNA-intercalating dye propidium iodide (PI). Whereas almost all untransfected cells or those transfected with an empty vector remained viable after 24 hr, many of those expressing Bim (*i.e.*, EE-antibody positive) contained sub-diploid DNA (Figure 4A). Indeed, by three days, 90% of the cells expressing Bim_L were dead (Figure 4B). The extent of cell death was proportional to the amount of *bim* DNA transfected (black bars, Figure 4C).

The cells expressing Bim appeared to die by apoptosis, as assessed by cell morphology and the generation of sub-diploid DNA (Figure 4A). As expected, the death process required activation of caspases, because co-expression of baculovirus p35, a competitive inhibitor of many types of caspases (Bump *et al.*, 1995), antagonised Bim-induced cell death, whereas an inactive mutant p35 did not (Figure 4C). Since crmA, a potent inhibitor of caspases 1 and 8 (ICE and FLICE) (Orth *et al.*, 1996; Srinivasula *et al.*, 1996) was not effective (Figure 4C), these particular caspases do not appear to play a critical role.

Numerous failed attempts to generate lines that stably express Bim suggested that it is toxic to diverse cell types. Those repeatedly tested include haemopoietic lines (FDC-P1, CH1, Jurkat, SKW6 and B6.2.16BW2), fibroblastoid lines (Rat-1, NIH3T3 and L929) and an epithelial line (293). The cells were electroporated with a vector encoding antibiotic resistance and either EE- or FLAG-tagged Bim_L and selected in antibiotic, but no line expressing Bim emerged. A vector encoding untagged Bim also failed to generate viable clones. We quantified the cytotoxicity of Bim by colony assays on transfected L929 fibroblasts. The EE-Bim_L vector yielded only one fifth as many antibiotic-resistant colonies as the control vector, and when six of the EE-Bim_L-transfected, drug-resistant colonies were expanded, only one contained any Bim and the level was very low (Table 1 and data not shown). Thus, high levels of Bim suppress clonogenicity and appear incompatible with prolonged cell viability.

- 62 -

EXAMPLE 4**Bim cytotoxicity can be abrogated by wild-type Bcl-2 but not inactive mutants**

Co-expression experiments established that Bcl-2 could block cell death induced by Bim_L (Figure 4D). In 293T cells transiently transfected with both the *bcl-2* and *bim_L* plasmids, relatively few cells died, even with a high concentration of *bim_L* DNA (compare the 4th sample in Figure 4C with the 3rd in Figure 4D). The cytotoxicity of *bim*, however, could not be countered by mutant forms of *bcl-2* rendered inactive by deletion of the BH4 homology region (Δ BH4) (Borner *et al.*, 1994), or by a point mutation in its BH1 (G145E) or BH2 (W188A) region (Yin *et al.*, 1994) (Figure 4D). Thus, ability to antagonise Bim-induced cell death required a functional Bcl-2 molecule.

High levels of Bcl-2 allowed stable expression of Bim_L. Indeed, when L929 cells stably expressing Bcl-2 were transfected with the EE-Bim_L vector, the frequency of antibiotic-resistant colonies approached that obtained with the control vector, and four of six colonies analysed contained moderate to high levels of Bim (Table 1 and data not shown). Similarly, using FDC-P1 clones expressing wt Bcl-2 (but not mutant Bcl-2), we could readily establish sub-clones expressing varying levels of Bim_L (Figure 5A). When grown in the presence of IL-3, all were indistinguishable in growth characteristics and morphology from the parental FDC-P1 cells or those bearing Bcl-2 alone. However, when deprived of IL-3 or irradiated, cells expressing Bcl-2 and a moderate or high level of Bim died more readily than those expressing Bcl-2 alone (Figure 5B). Since each clone had the same level of Bcl-2 (not shown), their sensitivity to apoptosis presumably reflects the ratio of the pro-apoptotic Bim to the anti-apoptotic Bcl-2.

- 63 -

EXAMPLE 5

The three isoforms of Bim all interact with Bcl-2 in vivo but vary in cytotoxicity

We next explored whether all isoforms of Bim were equivalent. An FDC-P1 clone
5 expressing human Bcl-2 was transfected with vectors expressing Bim_{EL}, Bim_L or Bim_S,
and puromycin-resistant clones that expressed the same amount of each isoform were
selected for further analysis (Figure 6A). To test for association with Bcl-2,
immunoprecipitates prepared from cell lysates using a monoclonal antibody specific for
human Bcl-2 were fractionated electrophoretically and blotted with anti-EE antibody.
10 Each of the Bim isoforms clearly bound to Bcl-2 (Figure 6B). However, when the
transfectants were deprived of IL-3 or subjected to γ -irradiation, it became evident that
Bim_S antagonised Bcl-2 more effectively than Bim_L while Bim_{EL} was the least potent
(Figures 6C). In addition, Bim_S suppressed L929 colony formation more effectively
than Bim_L or Bim_{EL} (Table 1). Thus, although all three Bim isoforms can bind to Bcl-2,
15 they vary in cytotoxicity, Bim_S being the most potent.

EXAMPLE 6

Bim binds to and antagonises Bcl-x_L and Bcl-w but not viral Bcl-2 homologues

20 To determine whether Bim interacts with other members of the Bcl-2 family, we
performed immunoprecipitation on lysates from 293T cells transiently co-transfected with
the relevant vectors. No interaction with the pro-apoptotic Bax protein was observed,
under conditions in which Bax:Bcl-x_L association was readily detectable. Association of
Bim with Bcl-x_L or each of three point mutants was assessed in ³⁵S-labelled 293T cells
25 (Figure 7A). Bim bound to wild-type Bcl-x_L but not to a mutant (mt 7) that lacks pro-
survival activity, nor to two mutants (mt 1 and mt 15) which retain significant
anti-apoptotic activity but cannot bind to Bax (Cheng *et al.*, 1996).

Bim_L also bound strongly to the other cellular pro-survival regulator tested, Bcl-w
30 (Gibson *et al.*, 1996) (Figure 7B). In marked contrast, Bim_L did not bind to either of

- 64 -

two virally encoded Bcl-2 homologues, the adenovirus E1B19K protein (Figure 7B) and the Epstein-Barr virus BHRF-1 protein, even though both viral proteins bound to EE-Bax. Thus, not all mediators of cell survival associate with Bim.

5 Functional tests mirrored the binding properties of the various Bcl-2 homologues. When transiently co-expressed with Bim in 293T cells, Bcl-x_L and Bcl-w countered Bim toxicity as effectively as Bcl-2 (Figures 7C and 7D). In contrast, little inhibition was observed with comparable levels of the mutant Bcl-x_L proteins (Figure 7C) or the adenovirus E1B19K protein (Figure 7D). These data suggest that Bcl-2-like inhibitors of apoptosis
10 must bind to Bim to inhibit its action.

EXAMPLE 7

The BH3 region is essential for interaction of Bim with Bcl-2 and for most of its ability to promote apoptosis

15 Since the BH3 region of several death-promoting proteins is essential for their activity (see Introduction), we tested a *bim*_L mutant lacking the BH3 region. In transfected cells the mutant protein (Δ BH3) was readily detected by immunofluorescence and Western blotting (Figure 8A), establishing that BH3 is not essential for stability of the
20 polypeptide. Unlike wt Bim, however, the Δ BH3 mutant did not bind to Bcl-2 *in vivo* (Figure 8B).

In some biological assays, the Δ BH3 mutant of Bim appeared inert. In contrast to wt Bim, it was easy to establish lines expressing Bim_L Δ BH3 from FDC-P1 (Figure 8A) or
25 L929 cells (Table 1). Moreover, Bim_L Δ BH3 did not impair the viability of the FDC-P1 cells in either the presence or absence of Bcl-2 (Figure 8C). Finally, 293T cells transiently transfected with Bim_L Δ BH3 exhibited high viability (not shown). These results indicate that the BH3 region is critical for Bim to promote apoptosis and suggest that Bcl-2 blocks this activity of Bim by binding to that domain. Importantly, however,
30 Bim_L Δ BH3 was not completely inactive. In the L929 clonogenicity assay, it still

- 65 -

markedly suppressed colony formation (Table 1). Thus, regions of Bim other than BH3 may promote apoptosis or interfere with clonogenicity in another way, such as by blocking cell growth.

5

EXAMPLE 8

Expression library screening and isolation of mouse and human bim cDNAs

Polyadenylated RNA prepared from p53^{-/-} KO52DA20 T lymphoma (Strasser *et al.*, 1994) cells subjected to γ -irradiation (10 Gy) was reverse-transcribed, using a
10 combination of oligo dT and random oligonucleotide primers, and ligated to EcoRI adaptors, using standard procedures. The cDNA was then ligated with Eco RI + Xho I-digested λ ZapExpress (Stratagene) arms and packaged *in vitro* according to the supplier's instructions. The resulting expression library was screened using radiolabelled Bcl-2 lacking the hydrophobic membrane localisation region. To prepare this probe,
15 cDNA encoding amino acids 1 to 210 of human Bcl-2 was subcloned into the vector pAR Δ R1 (Blanan and Rutter, 1992), and recombinant protein (FLAG-HMK-Bcl-2 Δ EC30) produced in IPTG-induced *E. coli* BL21pLysS (DE3) cells (Novagen) was purified on anti-FLAG M2 affinity gel (IBI Kodak) and then kinased *in vitro* using bovine heart muscle kinase (Sigma) and [γ -³²P]ATP (Amersham) (Blanan and Rutter, 1992). $\sim 10^6$
20 plaques were screened with $\sim 10^7$ cpm of the radiolabelled probe using the protocol of Blanan and Rutter (Blanan and Rutter, 1992). To reduce non-specific background, the filters were pre-incubated with lysates from induced parental BL21pLysS (DE3) cells and excess unlabelled ATP. Plaques that were positive on duplicate lifts were picked for two rounds of further screening. Positive clones were excised *in vivo* by coinfection with
25 filamentous ExAssist (Stratagene) helper phage and sequenced by automated sequencing (ABI Perkin Elmer). The human *bim* cDNA clone was isolated by screening human embryo and liver λ cDNA libraries (Stratagene) with an ~ 800 bp mouse *bim* cDNA probe, using standard techniques. The cDNAs were fully sequenced, analysed using Wisconsin GCG or DNASTAR software and compared with sequences in the Genbank
30 (including dBEST) and EMBL databases using the BLAST algorithm (Altschul *et al.*,

1990).

EXAMPLE 9

Expression constructs and site-directed mutagenesis

5

cDNAs were cloned into the expression vectors pEF PGKpuro (Huang *et al.*, 1997) or pEF PGKhygro (Huang *et al.*, 1997), or derivatives thereof incorporating N-terminal FLAG (DYKDDDDK) SEQ ID NO: 11 (Hopp *et al.*, 1988) or EE (EYMPME) SEQ ID NO: 12 (Grussenmeyer *et al.*, 1985) epitope tags. The *bim*ΔEBH3 mutation was
 10 generated by deleting the DNA encoding amino acids 94 to 100 (LRRIGDE) SEQ ID NO: 13 and replacing this with DNA corresponding to a Hind III site (encoding AL). Mutations in *bcl-2* (ΔBH4, G145E, W188A) (O'Reilly *et al.*, 1996; Huang *et al.*, 1997) were generated by polymerase chain reaction via splice overlap extension (Horton *et al.*, 1993) using the proof-reading *Pfu* DNA polymerase (Stratagene) (oligonucleotides used
 15 are detailed in SEQ ID NO: 14-26). The sequences of derived clones were verified by automated sequencing prior to function analysis.

EXAMPLE 10

Cell culture and transfection

20

Cell lines used were: mouse IL-3-dependent promyelocytic line FDC-P1; mouse T hybridoma B6.2.16BW2; mouse B lymphoma lines CH1 and WEHI 231; mouse pre-B lymphoma line WEHI 415 (derived from a tumour which arose in an Eμ-*myc* transgenic mouse); human B lymphoblastoid line SKW6; human T lymphoma line Jurkat; mouse T
 25 lymphoma lines WEHI 703, WEHI 707 (both derived from tumours which arose in Eμ-NRas transgenic mice) and WEHI 7.1; rat fibroblastoid line Rat-1; mouse fibroblastoid line NIH 3T3; mouse fibroblastoid line L929 subline LM(-TK); human embryonal kidney cell line 293 (ATCC CRL-1573) and SV40-transformed 293 cells, 293T (see Lithgow *et al.*, 1994; Strasser *et al.*, 1994; Strasser *et al.*, 1995; Huang *et al.*,
 30 1997). The procedures for culture and stable transfection are described elsewhere

- 67 -

(Huang *et al.*, 1997). Drug-resistant transfectants were cloned using the cell deposition unit of a FACStarPlus (Becton Dickinson) and clones expressing high levels of the protein of interest were identified by immunofluorescence staining of fixed and permeabilised cells followed by flow cytometric analysis.

5

EXAMPLE 11

Cell death assays

Cytokine deprivation and exposure to ionising radiation were the principal cell death
10 assays used to assess the sensitivity of FDC-P1 cells stably transfected with the various expression vectors. Cells were cultured in medium lacking cytokine or (in complete medium) after exposure to 10 Gy γ -radiation (provided by a ^{60}Co source at a rate of 3 Gy/min) and their viability determined over several days by vital dye (0.4% eosin) exclusion, as assessed by visual inspection in a hemocytometer, or by flow cytometric
15 analysis of cells that excluded propidium iodide ($5\mu\text{g/ml}$; Sigma) (Nicoletti *et al.*, 1991).

Cell death assays in 293T cells were performed after transient transfection of $\sim 5 \times 10^5$ cells using $6\mu\text{l}$ of Lipofectamine[®] (Gibco BRL) and a total of $1\mu\text{g}$ DNA in 2 ml of medium in 6 cm dishes; for co-transfections, *bim* plasmid (0.1, 0.2, $0.5\mu\text{g}$) was
20 co-transfected with $0.5\mu\text{g}$ of the other recombinant (eg *bcl-2*) plasmid and (0.4, 0.2, $0\mu\text{g}$) of empty vector. Forty-eight hours after transfection, the cells were harvested, fixed for 5 min in 80% methanol, permeabilised with 0.3% saponin (which was included in all the subsequent steps), and stained with $1\mu\text{g/ml}$ anti-EE monoclonal antibody (BabCO), followed by fluorescein isothiocyanate (FITC)-conjugated goat anti-mouse IgG ($1\mu\text{g/ml}$;
25 Southern Biotechnology) as the secondary agent and by $69\mu\text{M}$ propidium iodide in 38 mM sodium citrate pH 7.4 (Crissman *et al.*, 1990). Analysis was performed on a FACScan (Becton Dickinson), the proportion of dead cells being taken to be the proportion of EE-positive cells with less than 2C DNA content (Nicoletti *et al.*, 1991).

30 L929 fibroblast colony assays were performed in triplicate by scoring the numbers of

- 68 -

colonies in 10 cm dishes grown for 14-18 days with appropriate antibiotic selection. These cells had previously been split (1:3) from $\sim 10^6$ cells which had been transfected in 6 cm dishes 2 days earlier with 1 μ g total DNA and 12 μ l of Lipofectamine[®].

5

EXAMPLE 12**Immunofluorescence, immunoprecipitation and immunoblotting**

- Immunofluorescence staining of cytoplasmic proteins with the monoclonal antibodies Bcl-2-100 (mouse anti-human Bcl-2; [Pezzella *et al.*, 1990]) or mouse anti-EE (BabCO) followed by FITC-conjugated goat anti-mouse IgG (Southern Biotechnology) was performed as previously described (Huang *et al.*, 1997). Cells were analysed in the FACScan II (Becton Dickinson) after exclusion of dead cells on the basis of their forward and side scatter characteristics.
- 15 To investigate the subcellular localisation of EE-tagged Bim_L, transfected L929 fibroblasts grown in chamber slides (Erie Scientific Company, New Hampshire) were fixed in 4% paraformaldehyde for 10 min at room temperature and the slides were then allowed to dry and stored at -20°C. Prior to analysis by confocal microscopy, the cells were rehydrated by dipping the slides in water and then permeabilised for 15 minutes at room temperature in 0.5% Triton-X 100 in PBS. EE-Bim_L was detected by incubating for the cells with anti-EE monoclonal antibody for 30 minutes, washing several times in PBS containing 2% foetal calf serum and 0.05% Tween-20, and then incubating for 30 minutes with goat anti-mouse IgG conjugated to lissamine-rhodamine (Jackson Immunoresearch), all steps being performed at room temperature. Human Bcl-2 was detected similarly, using hamster anti-human Bcl-2 (6C8) (Veis *et al.*, 1993) followed by FITC-conjugated mouse anti-hamster IgG. Untransfected cells served as negative controls. Samples were analysed using a Leica confocal laser scanning microscope (Leica Lasertechnik).
- 25
- 30 To test for protein-protein interactions *in vivo*, immunoblotting was performed on stably

- 69 -

transfected FDC-P1 cells or transiently transfected 293T cells as described previously (Huang *et al.*, 1997). Briefly, lysates prepared from 10^5 - 10^6 cells were incubated with $\sim 5 \mu\text{g}$ antibody (anti-human Bcl-2, anti-FLAG M2 (IBI Kodak), or anti-EE monoclonal antibody), followed by protein G Sepharose (Pharmacia), and then pelleted, washed, 5 fractionated by SDS-PAGE and transferred to nitrocellulose membranes by electroblotting. The filters were incubated with mouse anti-human Bcl-2, anti-FLAG or anti-EE antibodies followed by affinity-purified rabbit anti-mouse IgG; bound antibodies were detected with ^{125}I -labelled staphylococcal protein A. In some experiments, the cells 10 were metabolically labelled with 100-200 $\mu\text{Ci/ml}$ of ^{35}S -methionine (NEG-072 from NEN) and equivalent TCA-precipitable counts (5×10^7 cpm) were used for each immunoprecipitation.

EXAMPLE 13

Antibody Production

15

Immunization

Wistar rats were immunized by injection of 100 μg of purified GST (glutathione-S-transferase)-Bim_L fusion protein, purified on a glutathione sepharose affinity matrix (Pharmacia, Uppsala, Sweden). For the first immunization the protein 20 was dissolved in complete Freund's adjuvant (Difco, Michigan, USA) and injected subcutaneously. Two subsequent boosts of the immunogen resuspended in incomplete Freund's adjuvant (Difco) were injected subcutaneously 3 and 6 weeks after the initial injection. A final boost was given four weeks later, i.e. three days prior to fusion.

25 *Cell Fusion and Hybridoma Culture*

Rat spleen cells were fused according to published procedures (Galfre *et al.*, 1977) with Sp2/0 mouse myeloma cells (Shulman *et al.*, 1978) at a ratio of 2:1 to 4:1, using polyethylene glycol 1500 (Boehringer Mannheim, Mannheim, Germany). After fusion, the cells were resuspended in Dulbecco's modified Eagle's medium (DMEM) containing 30 15% FCS (MultiSer, Trace Bioscience, Australia, batch #31104149), 5 mM

- 70 -

hypoxanthine, 0.02 mM aminopterin and 0.8 mM thymidine (HAT; Boehringer), with IL-6 and plated into flat bottom 96-well plates (Falcon, Becton Dickinson, NJ, USA). The source of IL-6 was supernatant from X63/0 hybridoma cells stably transfected with an IL-6 expression construct (Karasuyama *et al.*, 1988). The titre of IL-6 in the supernatant was determined by stimulation of the IL-6-dependent cell line 7TD1 (22). IL-6 was utilized in the fusion medium at a concentration which permitted maximal proliferation of 7TD1 cells. Fresh tissue culture medium was added to the hybridoma cells on day 7 after fusion and supernatants were harvested for analysis on days 9-11 depending on the rate of colony growth. Stable antibody-producing clones were established by two sequential steps of single cell cloning.

Tissue Culture and Cell Lines

The IL-3-dependent mouse promyelomonocytic cell line FDC-P1 (Dexter *et al.*, 1980) was cultured in the high glucose version of Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum (FCS), 50 μ M 2-mercaptoethanol, 13 μ M folic acid, 100 μ M L-asparagine and 1000 U/mL IL-3. The source of IL-3 was supernatant from X63/0 hybridoma cells stably transfected with an IL-3 expression construct (Karasuyama and Melchers, 1988). Derivative clones of FDC-P1 cells transfected with a human *bcl-2* expression construct, FDC-P1/Bcl-2 (Huang *et al.*, 1997), or a human *bcl-2* expression construct and a Glu-Glu (EE) epitope-tagged mouse *bim_L* expression construct, FDC-P1/Bcl-2/EE-Bim_L (O'Connor *et al.*, 1998) have been described previously.

Expression of Bim and Bcl-2 was verified by cytoplasmic immunofluorescent staining (see below and (Strasser *et al.*, 1995)) using 1 μ g/mL mouse anti-human Bcl-2 monoclonal antibody Bcl-2-100 (17) or 2 μ g/mL mouse anti-EE monoclonal antibody (anti-EE) (BabCO, Richmond, CA, USA) and as the secondary reagent 5 μ g/mL fluorescein isothiocyanate (FITC)-conjugated goat anti-mouse IgG antibodies (Southern Biotechnology, Birmingham, AL, USA).

Hybridoma screening

A 1:1 mixture of FDC-P1/Bcl-2 and FDC-P1/Bcl-2/EE Bim_L cells was prepared and fixed in 1% paraformaldehyde for 15 min at room temperature. Cells were washed twice in balanced salt solution containing 2% FCS and 10 mM sodium azide (wash buffer) and 5 x 10⁵ cells (in 50 µL wash buffer) were dispensed into each well of 96-well soft plastic U-bottomed plates (Dynex, VA, USA). The cell pellets were resuspended in 100 µL of day 10-11 hybridoma supernatant with 50 µL of a 1% saponin solution (for cell permeabilization) in wash buffer, vortexed and incubated for 30 min on ice. Plates were washed twice by centrifugation (3 min at 1500 rpm in a Heraeus Sepatech Megafuge 10 1.0R) in wash buffer containing 0.03% saponin and resuspended in 100 µL wash buffer containing 0.3% saponin and 10 µg/mL FITC-coupled goat anti-rat IgG (heavy and light chain-reactive) antibodies (Southern Biotechnology) and incubated for 30 min on ice. Finally, cells were washed twice in wash buffer containing 0.03% saponin and resuspended in wash buffer. To confirm Bim_L expression a control sample was stained 15 with mouse anti-EE antibodies (BabCO) and detected with 5 µg/mL goat anti-mouse IgG antibodies conjugated to FITC (Southern Biotechnology). Cell staining was analyzed in a FACScan (Becton Dickinson, Mountain View, CA, USA) after exclusion of dead cells on the basis of their forward and side light scatter characteristics. Fluorescence histograms were only printed from the positive samples. To speed up sample processing 20 to a rate of 250 to 300 samples per hour, only 500 cells were analyzed. This allowed one person to screen 2000 hybridoma cultures over two days.

Expression Constructs and Protein Purification

The vectors pEF Bcl-2 pGK_{puro}, pEF FLAG Bim_L and pEF EEBim_L pGK_{hygro} have 25 been described before (Huang *et al.*, 1997; O'Connor *et al.*, 1998). Full length mouse *bim_L* cDNA was cloned into pGEX (128/128) (Blancar and Rutter, 1992) to allow production of GST-FLAG-Bim_L protein in the bacterial strain JM109. The recombinant protein was purified from IPTG-induced (Sigma, St Louis, MO, USA) bacterial cultures using binding to glutathione-sepharose 4B and elution with reduced glutathione (Pharmacia) 30 (Smith and Johnson, 1988). The recombinant protein was resuspended in PBS pH 7.

- 72 -

Western blotting

FDC-P1/Bcl-2 and FDC-P1/Bcl-2/EE Bim_L cells were harvested in lysis buffer (20 mM Tris-HCl, pH 8.0, 125 mM NaCl, 1 mM EGTA, 1% Triton X-100, 10% glycerol, 0.5 μ g/mL Pefabloc, 1 μ g/mL of each: leupeptin, aprotinin, soybean trypsin inhibitor and pepstatin, 5 mM NaF and 2 mM Na₃VO₄, all reagents from Sigma or Boehringer Mannheim). Lysates from 10⁶ cells were boiled in gel running buffer (0.25 M Tris-HCl pH 6.8, 1% SDS, 20% glycerol, 5% 2-mercaptoethanol, 0.02% bromophenol blue), resolved on 4-20% polyacrylamide gels (Novex, San Diego, CA, USA) and transferred to nitrocellulose membranes by electroblotting. After incubation overnight at 4°C in 5% skimmed milk, 1% casein and 0.05% Tween-20 to prevent non-specific binding, the filters were incubated (1 hr at room temperature) with the rat anti-Bim monoclonal antibodies diluted 1:1 with blotting solution, followed by affinity-purified HRP-conjugated goat anti-rat IgG antibodies (Southern Biotechnology). Proteins were visualized by enhanced chemiluminescence (Amersham, Amersham, UK). Metabolic labelling of cells with ³⁵S-methionine and immunoprecipitation 293T human kidney embryonal cells were transiently transfected with Bim expression constructs, as described (O'Connor *et al.*, 1998). After 48 hours the cells were labeled overnight with 100 μ Ci/mL ³⁵S-methionine (Du Pont, NEN Research Products, Boston, MA, USA). Cell lysates were prepared in lysis buffer and quantified by TCA (trichloroacetic acid) precipitation. Equivalent TCA precipitable counts (10⁷ cpm) were used for each immunoprecipitation (O'Connor *et al.*, 1998), and analyzed on 4-20% gradient polyacrylamide gels (Novex). As a control anti-FLAG M2 (IBI Kodak, New Haven, CT, USA) and anti-mouse CD4 antibodies (clone H129.19.6.8) were used.

25

Results

Clones expressing high levels of Bcl-2 and EE-Bim_L were selected by continuous growth in the presence of both drugs and by immunofluorescent staining with monoclonal antibodies specific to Bcl-2 or the EE epitope tag (Figure 10A).

30

Rats were immunized with recombinant GST-Bim fusion protein. Hybridoma screening was performed 10-11 days after the fusion. The immunogen was tagged differently from the protein used for screening to avoid isolation of hybridomas producing tag-specific antibodies. A 1:1 ratio of FDC-P1/Bcl-2 and FDC-P1/Bcl-2/EE-Bim_L cells were fixed, permeabilized with saponin and used to screen hybridoma supernatants. Those culture wells containing antibodies to Bim produced a double fluorescence peak when analyzed by flow cytometry (Figure 10 D, E, F). The lower intensity peak represents background staining of FDC-P1/Bcl-2 cells. The higher fluorescence intensity peak is the result of specific Bim staining (Figure 10A). Culture supernatants with no Bim reactivity showed only a single peak of low fluorescence intensity (Figure 10B). Antibodies which were not specific to Bim but bound to some unknown epitope present in both FDC-P1/Bcl-2 and FDC-P1/Bcl-2/EE-Bim_L cells produced a single peak with high fluorescence intensity (Figure 10C). From the initial screen 18 potentially Bim reactive clones were expanded and sub-cloned. Three monoclonal antibodies were obtained, 4E4, 5E5 and 9F5, that stained Bim with high sensitivity and specificity (Figure 10 D-F).

Epitope mapping of monoclonal antibodies is possible with the technique described above, by using cell lines transfected with expression constructs encoding modified versions of the protein of interest. In the case of Bim we used FDC-P1/Bcl-2 lines that co-express different isoforms or mutants of Bim and found that the monoclonal antibodies detected only Bim_L.

Antibody isotyping was also achieved by a simple adaptation of the staining protocol (Figure 2). The mixture of FDC-P1/Bcl-2 cells and FDC-P1/Bcl-2/EE-Bim_L cells was fixed, permeabilized and stained with each of the three anti-Bim antibodies. As a secondary reagent biotinylated mouse anti-rat Ig isotype specific monoclonal antibodies were used: anti-rat IgG1 (RG11/39.4), anti-rat IgG2a (RG7/1.30) and anti-rat IgG2b (RG7/11.1) (Springer *et al.*, 1982) and in the final step FITC-coupled streptavidin. To determine the Ig light chain isotype of the antibodies, we used FITC-coupled Mar 18.5 (Lanier *et al.* 1982) mouse anti-rat Igκ antibodies. This analysis demonstrated that the

- 74 -

4E4 and 5E5 antibodies are IgG2b/ κ and the 9F5 antibody is IgG2a/ κ (Figure 11).

Bim could be detected in Western analysis by all three anti-Bim antibodies in lysates obtained from as few as 5×10^4 FDC-P1/Bcl-2/EE-Bim_L cells when binding of the
5 secondary antibody was revealed by enhanced chemiluminescence (ECL) (Figure 12). The three anti-Bim antibodies were also capable of immunoprecipitating Bim_L protein from 293T cells that had been transiently transfected with a FLAGbim_L expression construct (Figure 13). Both the Western blotting and immunoprecipitation assays clearly showed that the monoclonal anti-Bim antibodies were specific, as only Bim_L was seen in
10 FDC-P1/Bcl-2/EE-Bim_L cells and no non-specific protein was detected in FDC-P1/Bcl-2 cells.

EXAMPLE 14

Fine-Mapping of the Dynein Light Chain Binding Region in Bim

15 Bim_S, which lacks amino acids 42-71 found in murine Bim_L and 42-127 in murine Bim_{EL} (O'Connor, *et al*, EMBO J, 1998), is incapable of interacting with dynein light chain whereas a BH3 deletion mutant of Bim (Δ aa150-aa157) does bind. To determine precisely the amino acid residues required for binding, a fine mapping approach was
20 undertaken using the yeast-reverse-two-hybrid-system (Vidal, *et al*, PNAS, 1996). The system was set up so that binding of Bim_L (coupled to the GAL4 DNA binding domain) to dynein light chain (fused to the GAL4 activation domain) induces expression of orotidine-5-phosphate decarboxylase. This enzyme converts 5-fluoro-orotic acid (FOA) to 5-fluoro-uracil (5FU) which kills the cell. The region within Bim_L spanning amino
25 acids 1-149 was mutagenized by low fidelity PCR and recombined into the yeast vector encoding Bim_L. From 15,000 transformants 82 mutant clones of Bim_L which failed to interact with dynein light chain were selected in the presence of 5FOA. These clones were tested for full length Bim_L protein production by their ability to interact with Bcl-2. This was feasible because the BH3 region of Bim, necessary for interaction with Bcl-2, is
30 22 amino acids towards the N-terminus from the dynein light chain binding region

- 75 -

(O'Connor, *et al*, EMBO J, 1998). This led to the identification of 24 mutant clones of Bim_L which did not interact with dynein light chain but could still bind to Bcl-2. Residues D51, K52, S53, T54 and appeared to be most critical for binding of Bim_L to dynein light chain.

5

Four of the Bim_L mutants from the reverse yeast-two-hybrid screening were randomly chosen for further investigation: D51G, S53P, T54A and the double mutant T54I/N65S. A quantitative analysis of the strength of interaction between Bim_L or the four mutants with dynein light chain or, as a control Bcl-2, was performed in a yeast-two-hybrid assay
10 by measuring β -galactosidase activity using ONPG as the substrate. Wild-type Bim_L and the four Bim_L mutants had comparable strength of interaction with Bcl-2. In contrast, when binding to dynein light chain was studied it became apparent that S53P, T54A and T54I/N65S had less than 0.1 % of the activity compared to wild-type Bim_L while the D51G mutant was intermediary and retained approximately 5-10% of the affinity.

15

Interaction between wild-type Bim_L or the four Bim_L mutants with dynein light chain or Bcl-2 was also studied in co-immunoprecipitation assays. This was done in 293T cells transiently transfected with Bim, Bcl-2 and dynein light chain expression constructs and in FDC-P1 cells stably transfected with Bim and Bcl-2 expression constructs. These
20 experiments confirmed that all four mutants of Bim_L could efficiently interact with Bcl-2 but were unable to bind to dynein light chain. Collectively, these results demonstrate that amino acid residues D51, S53 and T54 within Bim are critical for interaction with dynein light chain.

EXAMPLE 15

25 Binding to Dynein Light Chain Regulates the Pro-Apoptotic Activity of Bim

Cytoplasmic dynein light chain is a component of the minus end directed dynein motor complex, an evolutionarily conserved microtubule bound ATPase, which is involved in flagellar movement in *Chlamydomonas* and retrograde organelle transport in mammalian
30 cells. Dynein heavy chain and dynein intermediate chains are integral structural

- 76 -

components of the dynein ATPase complex. Dynein light chain is also a stoichiometric component of this complex but its biochemical function is presently not known.

Bim_S, the splice variant of Bim, which does not bind to dynein light chain is a much more potent killer than Bim_L or Bim_{EL} (O'Connor *et al.*, EMBO J, 1998). FDC-P1 clones were generated which stably express Bcl-2 together with Bim_S, Bim_L or Bim_L bearing the mutations which abolish binding to dynein light chain and analysed their sensitivity to apoptotic stimuli. Three clones of each genotype, matched for equal levels of Bcl-2 and Bim proteins were selected for analysis. Parental FDC-P1 cells and 10 transfectants expressing only Bcl-2 were used as additional controls.

Upon cytokine deprivation or γ -irradiation FDC-P1 cells expressing Bcl-2 and Bim_S died much more rapidly than those expressing Bcl-2 plus Bim_L. Three of the Bim_L mutants, S53P, T54A and the double mutant T54I/N65S were as potent inducers of apoptosis as 15 Bim_S. In contrast, the D51G mutant, which retains some ability to bind to a dynein light chain, did not influence the killing potential of Bim and behaved like Bim_L. These results demonstrate that interaction with dynein light chain regulates the pro-apoptotic activity of Bim.

20

EXAMPLE 16

Immunohistochemical Analysis of Bim Expression in Mouse Tissues

Immunohistochemical staining intensity grading

0 negative

25 1+ weak positive

2+ moderately positive

3+ strongly positive

4+ extremely intense positive

30 *Salivary Glands*

- 77 -

Serous type cells of the parotid gland and the serous type cells of the submandibular gland (mixed serous and mucus secreting cells) 2-3⁺Bim immunoreactivity. The Mucus type cells of the sublingual gland had no Bim immunoreactivity.

5 *Pancreas*

Bim immunoreactivity (2-3⁺) detected on the ductal epithelial cells and more intensely (3-4⁺) in the cells of neuroendocrine origin in the islets. The exocrine/acinar tissue contained no Bim immunoreactivity (0).

10 *Thymus*

Thymic medulla strongly positive (3⁺), medullary thymocytes variably positive (0-3⁺). Cortical thymocytes mostly negative with scattered cells containing Bim immunoreactivity (0-3⁺).

15 *Spleen*

Megakaryocytes of the red pulp 3⁺ Bim immunoreactivity but RBC negative. B and T zone areas of the white pulp strongly Bim positive (3⁺), plasma cells also had strong Bim immunoreactivity.

20 *Kidney*

Weak Bim immunoreactivity detected on most tubular epithelia (1⁺), but strikingly more intense in the epithelial cells of the proximal convoluted tubules at the cortico-medullary junction (3-4⁺).

25 *Striated Muscle*

Appearance of a punctate pattern of Bim immunoreactivity along the muscle fibres (1-2⁺). When in the correct plain of section the transverse cross-strations of the cylindrical myofibrils in the sarcoplasm of the muscle fibres containing strong Bi, immunoreactivity (3⁺).

- 78 -

Liver

Liver hepatocytes and Kupffer cells have no Bim immunoreactivity (0), but the bile duct epithelial cells have moderate Bim immunoreactivity (2-3⁺).

5 *Intestine*

Small intestine: intense Bim staining detected in the enterocytes lining the villi (3-4⁺).

Colon: enterocytes of the shorter villi and also the cells lining the crypts and strong Bim immunoreactivity (3-4⁺).

10 *Heart*

Cross striations in the cytoplasm of cardiac involunatry striated muscle had prominent Bim immunoreactivity (3⁺), particularly around the outer walls of cardiac chambers and muscles around the outer walls of cardiac chambers and muscles around cardiac blood vessels.

15

Testes

Bim immunoreactivity was absent from sertoli cells Leydig cells, spermatogonia, spermocytes and spermatids (0), but mature sperm and residual bodies were strongly positive (4⁺).

20

Ovary

Follicular cells or ripening follicle prominently stained (2-3⁺), less intense Bim immunoreactivity observed in the interstitial cells, primordial follicles and corpus leteum (1-2⁺).

25

EXPRESSION ANALYSIS OF BIM IN CELL LINES BY IP WESTERN

	Cell line	Origin	Species	Bim Expression
	ALB 8.1	B lymphoma	mouse	+
	KO52 DA.20	T lymphoma	mouse	+
5	WEHI 703	T lymphoma	mouse	+
	B6.2.16.BW2	T lymphoma	mouse	++
	RAW 264.7	Macrophage	mouse	+
	J774/2	Macrophage	mouse	+
	F4N/3	erythroleukaemia	mouse	weak +
10	TS5	erythroleukaemia	mouse	weak +
	DP10	erythroleukaemia	mouse	weak +
	C2C12	muscle (myoblasts)	mouse	-
	L6	muscle (myoblasts)	rat	-
	416B	myeloid	mouse	+
15	P185X-2.1	mastocytoma	mouse	+
	FDC-P1	myeloid	mouse	-
	NIH3T3	fibroblast	mouse	+/-
	WEHI 11	sarcoma	mouse	+
	WEHI 164	sarcoma	mouse	-
20	S17	stromal	mouse	-
	L929	fibroblast	mouse	+/-
	MCF-7	breast carcinoma	human	++
	MDCK	kidney	dog	-
	293T	embryonic kidney	human	+
25	HK-2	kidney proximal	human	-
	G-401	Wilm's tumour	human	-
	TCMK-1	kidney	mouse	+
	Cosm6	kidney	monkey	+
	MH134	hepatoma	mouse	+
30	SW480	colon carcinoma	human	-
	EB-3	colon carcinoma	human	-

- 80 -

LIST OF RAT ANTI-BIM ANTIBODIES

	FACS			
CLONE	ISOTYPE	mBIM S	mBIM EL/L	huBIM EL/L
7H1	? K	+	+	-
8D1	? K	+	+	-
8F1	? K	+	+	-
9A12	IgG2aK	+	+	+
14A8	IgG2aK	+	+	+
16C4	IgG2aK	+	+	-
17C7	?	+	+	-
6A3	IgG1K	+	+	-
18D7	? K	+	-	-
5E8	?	+		
5E5	IgG2bK	-	+	+
4E4	IgG2bK	-	+	+
9F5	IgG2aK	-	+	+

LIST OF RAT ANTI-BIM ANTIBODIES

WESTERN			
mBIM S	mBIM L	mBIM EL	huBIM EL/L
+	+	+	+
+	+	+	-
+	+	+	-
+	+	+	+
+	+	+	+
weak	+	+	v. weak
weak	+	+	+
?	?	+	-
	+	+	+
-	+	? v. weak	+
-	+	? v. weak	+
-	+	? v. weak	+

EXAMPLE 17

Disruption of *Bim*

The *Bim* gene was inactivated by homologous recombination. The gene targeting vector (see Fig. 14) was assembled in *ploxPneo*-1 in which a neomycin phosphotransferase gene (*neo*^r), driven by a phosphoglycerate kinase (PGK) promoter, is flanked by bacteriophage P1 *loxP* sites. The 129/Sv mouse *Bim* genomic DNA sequences introduced at each end of the *loxP-neo*^r*loxP* cassette comprised the 4,000 bp region immediately upstream of the *Bim* exon 3 and the 2,700 bp downstream from exon 3. Introduction of a terminal herpes simplex virus thymidine kinase (*tk*) gene driven by a PGK promoter then completed the vector, which was linearized and electroporated into W9.5 ES cells (Koentgen et al., 1995). ES cell clones selected for resistance to G418 (i.e. *neo*^r gene integration) and gancyclovir (i.e. loss of the *tk* gene following homologous recombination) (Mansour et

- 82 -

al., 1988) were screened for homologous recombination at the *Bim* locus by Southern blot analysis. The *Bim* mutant ES cell clones were injected into the blastocoel cavity of 129/Sv blastocysts, which were then implanted into pseudopregnant foster mothers. Male chimeric progeny were crossed to 129/Sv females or, to delete the *neo^r* cassette, to B6/FVB F1 females expressing bacteriophage P1 Cre recombinase (Cre) (Laksoetal., 1996).

EXAMPLE 18

The genomic location of the mouse and human *Bim* genes

The genomic localisation of the mouse and human *Bim* genes was determined by hybridisation. ³H-thymidine labelled mouse *Bim_L* probes was used to probe a normal mouse metaphase spread. This demonstrated that the mouse *Bim* gene is located on chromosome 2 at bands F3-G. The corresponding localisation of human *Bim* gene was determined by fluorescence in situ hybridisation (FISH). A human *Bim_L* cDNA probe was nick-translated with biotin-14-dATP and hybridised in situ to normal male metaphases. The human gene is found in the syntenic region on chromosome 2 at bands 2q12-2q13.

EXAMPLE 19

Mutant mice lacking *Bim*

To determine the essential biological function of *Bim*, mice with a germline mutation in *Bim* have been generated. A number of characteristic abnormalities have been identified. Firstly, in an intercross of *Bim* +/- animals the number of *Bim* -/- offspring is significantly less than the expected 25%. This indicates that *Bim* may have an essential role in embryogenesis but that it can be partially compensated by related molecules. The *Bim* -/- mice that are generated have no obvious physical abnormality and both females and males are fertile. It is possible that the genetic background influences the phenotype of the *Bim* -/- mutation. Analysis of the haematopoietic compartment has shown that the

- 83 -

Bim ^{-/-} mice have increased numbers of blood leukocytes (~2-fold) and increased numbers of spleen cells (also ~2-fold). In the thymus the numbers of CD4⁺CD8⁺ pre T cells are reduced (~2-fold) and the numbers of mature CD4⁺CD8⁻ and CD4⁺CD8⁺ T cells are increased (~2- to 3-fold). Cell survival analysis on purified CD4⁺CD8⁺ pre T cells demonstrated that the *Bim* ^{-/-} cells are considerably more resistant to a range of apoptotic stimuli (growth factor deprivation, corticosteroids, DNA damage, calcium ionophores and phorbol esters) compared to control cells. The cells from the heterozygous *Bim* ^{+/-} mice were also more resistant than the cells from normal mice.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

BIBLIOGRAPHY

Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990) *J. Mol. Biol.*, **215**, 403-410.

Blancar, M. and Rutter, W. (1992) *Science* **256**, 1014-1018 .

Boise, L. H., Gonzalez-Garcia, M., Postema, C. E., Ding, L., Lindsten, T., Turka, L. A., Mao, X., Nunez, G. and Thompson, C. B. (1993) *Cell* **74**, 597-608.

Boise, L. H., Minn, A. J., Noel, P. J., June, C. H., Accavitti, M. A., Lindsten, T. and Thompson, C. B. (1995) *Immunity* **3**, 87-98.

Borner, C., Martinou, I., Mattmann, C., Irmeler, M., Schaerer, E., Martinou, J.-C. and Tschopp, J. (1994) *J. Cell Biol.*, **126**, 1059-1068.

Boyd, J. M., Gallo, G. J., Elangovan, B., Houghton, A. B., Malstrom, S., Avery, B. J., Ebb, R.G., Subramanian, T., Chittenden, T., Lutz, R. J. and Chinnadurai, G. (1995) *Oncogene*, **11**, 1921-1928.

Cheng, E. H.-Y., Levine, B., Boise, L. H., Thompson, C. G. and Hardwick, J. M. (1996) *Nature*, **379**, 554-556.

Chittenden, T., Flemington, C., Houghton, A. B., Ebb, R. G., Gallo, G. J., Elangovan, B., Chinnadurai, G. and Lutz, R. J. (1995) *EMBO J.*, **14**, 5589-5596.

Chittenden, T., Harrington, E. A., O'Connor, R., Flemington, C., Lutz, R. J., Evan, G. I. and Guild, B. C. (1995) *Nature*, **374**, 733-736.

Crissman, H. A., Darzynkiewicz, Z., Steinkamp, J. A. and Tobey, R. A. (1990) *Methods Cell. Biol.*, **33**, 305-314.

Dexter, T. M., G. D. Scott, E. Scolnick, and D. Metcalf (1980) *J. Exp. Med.* **152**: 1036-1047.

Farrow, S. N., White, J. H. M., Martinou, I., Raven, T., Pun, K.-T., Grinham, C. J., Martinou, J. C. and Brown, R. (1995) *Nature*, **374**, 731-733.

Galfre, G., S. C. Howe, C. Milstein, G. W. Butcher, and J. C. Howard (1977) *Nature* **266**: 550-552.

Gibson, L., Holmgreen, S., Huang, D. C. S., Bernard, O., Copeland, N. G., Jenkins, N. A., Sutherland, G. R., Baker, E., Adams, J. M. and Cory, S. (1996) *Oncogene*, **13**, 665-675.

Grussenmeyer, T., Scheidtmann, K. H., Hutchinson, M. A., Eckhart, W. and Walter, G. (1985) *Proc. Natl. Acad. Sci. USA*, **82**, 7952-7954.

Han, J., Sabbatini, P. and White, E. (1996) *Mol. Cell. Biol.*, **16**, 5857-5864.

Hanada, M., Aime-Sempe, C., Sato, T. and Reed, J. C. (1995) *J. Biol. Chem.*, **270**, 11962-11969.

Henderson, S., Huen, D., Rowe, M., Dawson, C., Johnson, G. and Rickinson, A. (1993) *Proc. Natl. Acad. Sci. USA*, **90**, 8479-8483.

Horton, R. M., Ho, S. N., Pullen, J. K., Hunt, H. D., Cai, Z. and Pease, L. R. (1993) *Methods Enzymol.*, **217**, 270-279.

- 86 -

- Huang, D. C. S., Cory, S. and Strasser, A. (1997) *Oncogene*, **14**, 405-414.
- Huang, D. C. S., O'Reilly, L. A., Strasser, A. and Cory, S. (1997) *EMBO J.*, **16**, 4628-4638.
- Huang, D. C. S., S. Cory, and A. Strasser (1997) *Oncogene* **14**: 405-414.
- Inohara, N., Ding, L., Chen, S. and Núñez, G. (1997) *EMBO J.*, **16**, 1686-1694.
- Jacobson, M. D. (1997) *Curr. Biol.*, **7**, R277-R281.
- Jacobson, M. D., Weil, M. and Raff, M. C. (1997) *Cell*, **88**, 347-354.
- Karasuyama, H., and F. Melchers (1988) *Eur. J. Immunol.* **18**: 97-104.
- Kerr, J. F. R., Wyllie, A. H. and Currie, A. R. (1972) *Br. J. Cancer*, **26**, 239-257.
- Kiefer, M. C., Brauer, M. J., Powers, V. C., Wu, J. J., Umansky, S. R., Tomei, L. D. and Barr, P. J. (1995) *Nature*, **374**, 736-739.
- Koentgen *et al.* (1995). *Genes Develop* **9**: 1965-1977
- Krajewski, S., Tanaka, S., Takayama, S., Schibler, M. J., Fenton, W. and Reed, J. C. (1993) *Cancer Res.*, **53**, 4701-4714.
- Kroemer, G. (1997) *Nature Med.*, **3**, 614-620.
- Lakso *et al.* (1996). *Proc. Natl. Acad. Sci. USA* **93**: 5860-5865
- Lanier, L. I., G. A. Gutman, D. E. Lewis, S. T. Griswold, and N. L. Warner (1982) *Hybridoma* **1**: 125-131.

Lithgow, T., van Driel, R., Bertram, J. F. and Strasser, A. (1994) *Cell Growth Differ.*, **5**, 411-417.

Mansour *et al.* (1988). *Nature* **336**: 348-352

Monaghan, P., Robertson, D., Amos, T. A. S., Dyer, M. J. S., Mason, D. Y. and Greaves, M. F. (1992) *J. Histochem. Cytochem.*, **40**, 1819-1825.

Muchmore, S. W., Sattler, M., Liang, H., Meadows, R. P., Harlan, J. E., Yoon, H. S., Nettesheim, D., Chang, B. S., Thompson, C. B., Wong, S.-L., Ng, S.-C. and Fesik, S. W. (1996) *Nature*, **381**, 335-341.

Nicholson, D. W. and Thornberry, N. A. (1997) *Trends Biochem. Sci.*, **22**, 299-306.

Nicoletti, I., Migliorati, G., Pagliacci, M. C., Grignani, F. and Riccardi, C. (1991) *J. Immunol. Meth.*, **139**, 271-279.

O'Connor, L., A. Strasser, L. A. O'Reilly, G. Hausmann, J. M. Adams, S. Cory, and D. C. S. Huang (1998) *EMBO J.* **17**: 384-395.

O'Reilly, L., Huang, D. C. S. and Strasser, A. (1996) *EMBO J.*, **15**, 6979-6990.

Oltvai, Z. N., Millman, C. L. and Korsmeyer, S. J. (1993) *Cell*, **74**, 609-619.

Pezzella, F., Tse, A. G. D., Cordell, J. L., Pulford, K. A. F., Gatter, K. C. and Mason, D. Y. (1990) *Am. J. Path.*, **137**, 225-232.

Reed, J. C. (1997) *Nature*, **387**, 773-776.

Sattler, M., Liang, H., Nettesheim, D., Meadows, R. P., Harlan, J. E., Eberstadt, M.,

- Yoon, H. S., Shuker, S. B., Chang, B. S., Minn, A. J., Thompson, C. B. and Fesik, S. W. (1997) *Science*, **275**, 983-986.
- Sedlak, T. W., Oltvai, Z. N., Yang, E., Wang, K., Boise, L. H., Thompson, C. B. and Korsmeyer, S. J. (1995) *Proc. Natl. Acad. Sci. USA*, **92**, 7834-7838.
- Shulman, M., C. D. Wilde, and G. Köhler (1978) *Nature* **276**: 269-270.
- Smith, D. B., and K. S. Johnson (1988) *Gene* **6788**: 31-40.
- Springer, T. A., A. Bhattacharya, J. T. Cardoza, and F. Sanchez-Madrid (1982) *Hybridoma* **1**: 257-273.
- Strasser, A., Harris, A. W., Huang, D. C. S., Krammer, P. H. and Cory, S. (1995) *EMBO J.*, **14**, 6136-6147.
- Strasser, A., Harris, A. W., Jacks, T. and Cory, S. (1994) *Cell*, **79**, 329-339.
- Vaux, D. L., Cory, S. and Adams, J. M. (1988) *Nature*, **335**, 440-442.
- Veis, D. J., Sentman, C. L., Bach, E. A. and Korsmeyer, S. J. (1993) *J. Immunol.*, **151**, 2546-2554.
- Wang, K., Yin, X.-M., Chao, D. T., Milliman, C. L. and Korsmeyer, S. J. (1996) *Genes Dev.*, **10**, 2859-2869.
- White, E., Sabbatini, P., Debbas, M., Wold, W. S. M., Kusher, D. I. and Gooding, L. R. (1992) *Mol. Cell. Biol.*, **12**, 2570-2580.
- Yin, X.-M., Oltvai, Z. N. and Korsmeyer, S. J. (1994) *Nature*, **369**, 321-323.

- 89 -

Zha, H., Aimé-Sempé, C., Sato, T. and Reed, J. C. (1996) *J. Biol. Chem.*, **271**, 7440-7444.

End
B2 B2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
22